

2014 WHEAT CRP ANNUAL REPORT

for the Consortium and the Fund Council





Research Program on WHEAT

Contents

Acronyms and abbreviations	iv
Part I – Technical report	1
A. Key messages	1
A.1 Progress and challenges in 2014	1
A.2 Synthesis of two most significant achievements	1
A.3 Financial summary	2
A.4 Gender	2
A.5 CRP governance and management Improvements	3
B. Impact pathway and intermediate development outcomes (IDOs)	3
C. Progress along the impact pathway	3
C.1 Progress towards outputs	3
C.2 Progress towards the achievement of outcomes	5
C.3 Progress towards impact	6
D. Gender research achievements	7
D.1 Selected gender research achievements	7
D.2 Success and challenges in mainstreaming gender research	7
E. Partnership building achievements	7
E.1 Selected partnership building achievements	7
E.2 Strategic partnerships	8
E.3 Interactions with other CRPs	8
F. Capacity building	9
G. Risk management	9
H. Lessons learned	10
H.1 Level of confidence of the response to the key performance indicators	10
H.2 Unintended results and innovative initiatives	10
Part II: Financial report	11
Annex 1. Indicators of progress in 2014	16
Annex 2: Performance indicators for gender mainstreaming with targets defined	20
Annex 3. How WHEAT delivers towards impacts (CGIAR System Level Outcomes)	21
Annex 4. WHEAT-related impact studies published during 2004-2014	22
Annex 5. 2014 WHEAT publications	23

Acronyms and abbreviations

<u>A4NH</u>	CGIAR Research Program on Agriculture
	for Nutrition and Health
<u>ACIAR</u>	Australian Centre for International
	Agricultural Research
APR	Adult plant resistance
<u>AgMIP</u>	Agricultural Model Intercomparison and
	Improvement Project
<u>AIP</u>	Agricultural Innovation Program for
	Pakistan
AR4D	Agricultural research for development
ATA	Nigerian Government's Agricultural
	Transformation Agenda
<u>AVRDC</u>	World Vegetable Center
BBSRC	UK Biotechnology and Biological Sciences
	Research Council
<u>BGRI</u>	Borlaug Global Rust Initiative
<u>BMGF</u>	Bill & Melinda Gates Foundation
<u>BMZ</u>	Federal Ministry for Economic
	Cooperation and Development, Germany
CA	Conservation agriculture
<u>CAAS</u>	Chinese Academy of Agricultural Sciences
<u>CCAFS</u>	CGIAR research program on Climate
	Change, Agriculture and Food Security
<u>CCAP</u>	Center for Chinese Agricultural Policy
<u>CIAT</u>	International Center for Tropical
	Agriculture
CPG	WHEAT competitive partner grant
<u>CSIRO</u>	Australia's Commonwealth Scientific and
00104	Industrial Research Organization
<u>CSISA</u>	Cereal Systems Initiative for South Asia
DAriseq	Diversity Arrays Technology
	Durable Rust Resistance in Wheat Project
EIAR	Ethiopian institute of Agricultural
	Research
FACASI	Agriculture for Sustainable Intensification
	Agriculture for Sustainable Intensincation
FARA	Forum for Agricultural Research in Africa
FD	Flagshin Projects
Fe	Iron
GBS	Genotyping-by-sequencing
GMO	Genetically modified organism
GRiSP	Global Rice Science Partnership
GS	Genomic selection
ha	hectares
HeDWIC	Heat and Drought Wheat Improvement
	Consortium (HeDWIC)
ICAR	Indian Council of Agricultural Research

ICRISAT	International Crops Research Institute for
	the Semi-Arid Tropics
	Intermediate development outcomes

- IDOs Intermediate development outcomes
- ILRI International Livestock Research Institute
- IRRI International Rice Research Institute IWIN International wheat improvement
- network
- IWWIP International Winter Wheat Improvement Program
 - IWYP International Wheat Yield Partnership
- JIRCAS Japan International Research Center for Agricultural Sciences
- <u>MasAgro</u> Modernización Sustentable de la Agricultura Tradicional
 - MC Management committee
- MEXPLAT Mexican Phenotyping Platform <u>MNFSR</u> Pakistan Ministry of National Food Security and Research
 - MoA Ministry of Agriculture
 - Mt Million tons
 - N Nitrogen
 - NARS National agricultural research systems
 - PARC Pakistan Agricultural Research Council
 - PLL Precision laser land leveling
 - QTL Quantitative trait loci
 - R&D Research and development
 - <u>RWC</u> Rice Wheat Consortium for Indo-Gangetic Plains
 - RSs WHEAT Research Strategies
- SARD-SC Support to Agricultural Research for Development on Strategic Commodities of the African Development Bank
 - Seeds of Discovery
 - SIAP Mexico's Servicio de Información Agroalimentaria y Pesquera
 - SLO System level outcomes
 - t tons
- UC-Davis University of California, Davis USAID U.S. Agency for International
 - Development
 - W4A Wheat for Africa
 - WIT Jeanie Borlaug Laube Women in Triticum Early Career Award
 - W-ISC WHEAT-Independent Steering Committee
 - WYC Wheat Yield Consortium
 - WYCYT International Wheat Yield Consortium Yield Trial
 - YR Yellow rust
 - Zn Zinc
 - ZT Zero tillage

Part I – Technical report

A. Key messages

A.1 Progress and challenges in 2014

In the spring of 2014, India, Pakistan, CIMMYT and WHEAT celebrated the 100th birthday of Dr. Norman Borlaug, drawing attention to present and future challenges to wheat production. High on the list are expanding demand and rising temperatures: A study by WHEAT scientists published in *Nature Climate Change* has shown that, globally, for every 1° C increase in growing season mean temperatures, wheat production decreases by 6% — a worldwide loss of 42 million tons of grain — and rising temperatures are already reducing global wheat production (see p.3). Regionally, another WHEAT study has documented a slowdown in average wheat cultivar turnover rates in India from 9 years (1997) to 12 years (2009; see p.3), increasing risks and highlighting the challenge that WHEAT and its national partners face in many world regions. Later in the year, the <u>World Food Prize</u> was awarded to distinguished wheat breeder <u>Dr. Sanjaya Rajaram</u>, who warned that "future crop production is bound to decline unless we fully factor in the issues related to climate change, soil fertility and water deficits. It will require all the resources of international research centers, national governments, foundations, NGOs and farmer groups together to synergize future agricultural technologies and food production," underlining the importance of partnerships.

As wheat agriculture and food security challenges loom large, research funded by the WHEAT Partner Budget documented the past impact of germplasm and training in China, the world's largest wheat producer and, together with India, home to half of the world's poor (See: Where Do the World's Poor Live? A New Update, Andy Sumner, IDS Working Paper 393, Institute of Development Studies, 2012, ISBN: 978-1-78118-061-7). WHEAT researchers documented the long-term productivity impacts of agronomic technology adoption by farmers in the Indo-Ganges Plain, guiding future investments in that region. The WHEAT Global Variety Release and Adoption study will near completion in mid-2015.

With regard to outputs, sustainable intensification successes were achieved in Mexico, South Asia and Ethiopia, whilst improved wheat varieties development, distribution and adoption saw progress Afghanistan, Central Asia, South Asia and Sub-Saharan Africa (p.6), partly in collaboration with other CRPs (A4NH, p.6).

Bilateral and Windows 1&2 investments supported expanded capacity for gender research and mainstreaming (p.7), with greater gender equity achieved in capacity development (p.9). Last but not least, WHEAT developed an excellent Extension Period Proposal, which guided the reorganization of WHEAT research domains (five Flagship Projects) and changes to CRP Governance and Management initiated in 2014.

A.2 Synthesis of two most significant achievements

Impacts in China of international wheat partnerships. A 2014 study¹ led by the Director of the Center for Chinese Agricultural Policy (CCAP) of the Chinese Academy of Sciences shows that Chinese breeders' use of CIMMYT germplasm increased the country's wheat total factor productivity as much as 14% in the past three decades (annual TFP growth rate of between 0.17% and 0.45%), representing as much as 10.7 million tons of added grain worth US \$3.4 billion, based on 2011 prices. Results also showed that breeders' use of CIMMYT germplasm has been increasing: CIMMYT contributions are present in more than 26% of all major wheat varieties released in China after 2000 and have added valuable diversity for

¹ J. Huang, C. Xiang and Y. Wang. 2015. <u>The Impact of CIMMYT Wheat Germplasm on Wheat Productivity in China.</u> Mexico, D.F.: CGIAR Research Program on Wheat.

important traits, including yield, protein content, disease resistance and earlier maturity. A separate, three-year field experiment in the Sichuan Basin² showed that cultivars developed from CIMMYT synthetic hexaploid wheats yielded on average 11.5% more than local, elite non-synthetic-derived cultivars. Another study by CCAP showed that 350 Chinese researchers (15% female during 1980-1990; rising to 35% female during 2000-2012 and since the 1990's increasingly focused on younger scientists) had taken part in CIMMYT wheat training programs since 1970. Of those, 170 benefited from visiting scientist appointments at CIMMYT; many of the alumni now hold important positions in China's wheat research system.

Water-saving impacts in the Indo-Gangetic Plain: Long-term investments pay off. A 2014 overview paper³ extensively evaluated leading agronomic recommendations for South Asia. Precision laser land leveling (PLL) is the most effective innovation for reduced and equitable usage of inputs like water and maximum production, especially in coarse- and medium-textured soils. Study and promotion of the technique expanded through the Rice-Wheat Consortium during 1994-2008. It is now practiced on over 1.5 million hectares in South Asia and allows farmers to save 15-30% irrigation water and benefit from up to 6% higher yields in rice, wheat and other crops, compared to traditionally-leveled fields. Reports from India have documented wheat yields as much as 16.6% higher, with nearly 50% less irrigation water use and 132% higher water productivity with PLL and raised bed planting, compared to traditional practices. Use of laser leveling in Pakistan started in 1985 through an on-farm water management program in Punjab; in a 2013 report the technology was shown to save 31% irrigation water, raise water productivity to 1.58 kg/m³ and increase fertilizer use efficiency by nearly 4%.⁴

\$M	As per PIA	Budget	Actual Spend	of which on Gender
W1&2*	14.327	19.54	18.19	4.92; of total 2014 spend
W3	Part of bilateral figure below	13.7	9.43	See above
Bilateral	25.505	17.46	13.57	See above

A.3 Financial summary

The 2014 budget and expenditures incorporate WHEAT budget and ICARDA decentralization funding. The latter is not considered in the PIA signed in 2011. The actual Consortium Board approved budget was US \$ 13.9 M; by late Nov 2014 it was US \$16.6 M. This was due to increased W2 income, now half of WHEAT's combined W1&2 income. See the *Financial Report* (p.11), for more detail.

A.4 Gender

Gender performance self-assessment as per Annex 2, p.20, is "approaching requirements."

In 2014 WHEAT has increased investments in strategic gender research, in-house gender capacity building and making a growing number of projects gender responsive. The main thrust of strategic gender research is the cross-CRP "Global Study on Gender Norms, Agency and Innovation Study in Agriculture and Natural Resource Management" (Gennovate). This is particularly important for WHEAT,

² <u>https://www.agronomy.org/publications/cs/abstracts/55/1/98</u>.

³ <u>http://www.sciencedirect.com/science/article/pii/B9780128001318000042</u>.

⁴⁴ http://www.uet.edu.pk/research/researchinfo/13-RJ-JUL-2013/2-ART-7.pdf.

as Gennovate will improve on the very limited evidence base regarding gender and wheat. WHEAT gender experts co-lead this work with colleagues from 12 other CRPs. The CIMMYT-led CRPs WHEAT and MAIZE have already completed data collection for more than 19 of the 70+ case studies planned. WHEAT is also working with gender experts across other CRPs to secure funding for Gennovate's critical analysis and write-up phase.

A.5 CRP governance and management improvements

WHEAT-Independent Steering Committee (W-ISC) members elected an Independent Chair. W-ISC membership expanded to eight voting members, as well as ex-officio, non-voting members CIMMYT and ICARDA DGs and Chairs of their Boards' Program Committee. A CRP Director has been appointed, starting 1st January 2015. For details, see this <u>note on WHEAT management and governance changes</u>.

B. Impact pathway and intermediate development outcomes (IDOs)

Following a standardization of CRP structures, the WHEAT strategy was reorganized around 5 Flagship Projects (FPs) in 2014, encompassing the 10 Strategic Initiatives of the original WHEAT proposal (see Annex 3, Figure 2). The FPs implement the two WHEAT Research Strategies (RSs) and contribute to a number of Intermediate Development Outcomes (IDOs) and, as part of the CRP Portfolio, to the three System Level Outcomes (SLOs) (see Annex 3, Figure 3). Each FP comprises from three to six Clusters of Activities (CoAs), which represent aggregations of projects and work packages within a research thrust. WHEAT contributes to the following IDOs: IDO 1-Productivity, IDO 2-Food Security, IDO 4-Income, IDO 5 – Gender and Empowerment, IDO 6-Capacity to Innovate and IDO 9-Environment. The CRP also contributes to the other IDOs through joint efforts with other CRPs.

The completion rate on 2014 deliverables for projects under WHEAT ranges from 83 to 100%, with an overall output achievement above 97%, as documented in WHEAT reporting templates (for W1&2) and bilateral progress reporting (see Annex 1, p. 16 for details). An overview of FP outputs delivering towards sub-IDOs and SLOs is provided at Annex 3, p. 21. A list of impact studies published during 2004-2014 and their key findings appears in Annex 4, p. 22.

C. Progress along the impact pathway

C.1 Progress towards outputs

Rising Temperatures, Falling Wheat Yields. FP 1. A 2014 paper⁵ published in *Nature Climate Change*, discusses the work of 50 scientists (including 2 WHEAT) from 15 countries who used data from 30 crop models and widely distributed field experiments to prove that, for every 1° C increase in growing season mean temperatures, wheat production decreases by 6% — equivalent to a worldwide loss of 42 million tons of grain — and that rising temperatures are already reducing global wheat production. The results link to other ex-ante research under WHEAT and CCAFS and drive work like that of the Heat and Drought Wheat Improvement Consortium (HeDWIC; see more under "partnerships").

Varietal turnover: Why do farmers grow old varieties? FP 1. A 2014 study⁶, led by a professor at the University of Göttingen, Germany and former CIMMYT socioeconomist working in the Cereal Systems Initiative for South Asia (<u>CSISA</u>) program, explains how and why cultivar depreciation accentuates the vulnerability of resource-poor farmers to production risks, showing that average wheat cultivar turnover rates in India slowed from 9 years in 1997 to 12 years in 2009.

⁵ <u>Rising temperatures reduce global wheat production</u>. (If you have trouble, right-click on the title, copy the link, and paste it into your browser.)

⁶ In press with the *Journal of Agricultural Science*, and available as an IFPRI discussion paper.

More and better markers for key traits and genotypic analysis of genetic diversity. FP 2. We are switching technologies, trying to move from gel electrophoresis to fluorescence-based genotyping systems to reduce the time and cost. So instead of running SSRs we will run KASP assays and promote the KASP system, which works well for wheat, especially for the gene-based markers. It will take two-to-three years to move completely to this system; right now about 50-60% of our data are KASP SNPs The technology is being used both in CIMMYT and ICARDA labs, in collaboration with service providers.

Pre-Breeding for yield potential: Data from 2nd WYCYT and performance of pipeline material. FP 2.

Using information on photosynthetic and partitioning traits, hybridization schemes were designed to combine physiological traits and achieve cumulative gene action for yield potential. The resulting germplasm has produced superior yields and biomass in global, multi-location trials (the International Wheat Yield Consortium Yield Trial – WYCYT; and preliminary results from the Stress Adaptive Trait Yield Nursery -- SATYN). The top-yielding WYCYT line has a Mexican landrace in its pedigree and pipeline materials are dominated by synthetic derivatives and landraces in their pedigrees.⁷

Precision phenotyping takes off. FP 3. Indices derived from airborne imagery (drone- or blimp-mounted multi-spectral cameras or other sensors) of breeding plots were validated against equivalent ground-based indices (correlations between 0.58 and 0.91) for thousands of breeding lines, and provided significant, better-than-equivalent correlations with yield. The approach will be refined for field-based screening of large collections of genetic resources to identify new sources of adaptive traits, in gene discovery and in mainstream breeding for quantitative traits.

Global partnerships for phenotyping. FP 3. The Izmir, Turkey, phenotyping platform for yellow rust was initiated in 2012 and field facilities are operating; the aim is to complete the laboratory and containment facilities in 2015. The Mexican Phenotyping Platform (MEXPLAT) was established and is running as core infrastructure, with collaborative research and training and a focus on understanding yield potential and related processes. An agreement was signed with national partners in Sudan (early, terminal heat stress), Tunisia (Septoria, leaf rust) and Uruguay (multiple biotic stresses) to set up three additional precision phenotyping platforms (total target: 15 platforms) across the globe. During 2015-16 and with selected national partners, WHEAT will aim to establish more wheat phenotyping platforms, depending on W1&2 income and NARS co-investment, to expand the precision and predictive value of phenotyping data for new germplasm from WHEAT and partners' breeding pipelines.

More durable resistance to wheat diseases. FP 3. CIMMYT and ICARDA are deploying varieties that possess adult plant resistance (APR) based on combinations of minor, slow rusting genes that provide high levels of more durable resistance. Four characterized genes have pleiotropic effects, conferring partial APR to all three rusts and powdery mildew. High-yielding, rust resistant lines generated through Mexico-Kenya shuttle breeding are being distributed in CIMMYT international yield trials and screening nurseries. Gene cloning and gene function descriptions and other work were done at Australia's Commonwealth Scientific and Industrial Research Organisation (CSIRO) the University of Zurich, Agriculture and Agri-Food Canada, Iowa State University, the Plant and Food Research-New Zealand.⁸ Results from discovery (upstream) research under FP2 are moving to breeding in FP3, guided by ex-ante impact studies under FP1. Investments in upstream R&D aim to assure future success in past and current downstream R4D on rust. W1&2 funded research complements a bilaterally-funded project (DRRW/Cornell) and bilaterally-funded impact studies by a US partner (USAID Linkage/<u>University of Minnesota</u>)⁹ to guide targeting of WHEAT investments under FP2&3. ICARDA is combining 2-3 major

⁷ <u>Reynolds et al. (eds.) 2015</u>, pp. 17-18.

⁸ See <u>doi: 10.3389/fpls.2014.00641</u> and <u>doi: 10.1007/s00122 -014-2390-z</u>.

⁹ See <u>doi: 10.1126/science.1229707</u>.

genes in minor gene background, using molecular markers and inter-country shuttle followed by yield testing at key locations, shuttling of the segregating generation between Lebanon and Ethiopia, followed by key location testing in Lebanon, Morocco, Egypt, Sudan and Ethiopia.

Nutritious wheat. FP 3. Some of the high-zinc lines developed under HarvestPlus (A4NH) are good performers under drought and late sown heat conditions and have become part of the global breeding platform (FP3.3). After nearly 10 years of intensive breeding and selection for Zn and Fe traits, trials have been released in over 10,000 farmers' fields in India and are expected to be released to farmers in Pakistan in early 2015. As recent evidence shows that grain zinc and/or iron is controlled by a few major QTLs and a few minor QTLs, with moderate to high heritability, WHEAT is going to integrate breeding for high Zn and Fe into regular breeding programs. See also Section E.3.

Conservation agriculture for wheat in Asia, North Africa, and the Middle East. FP 4. Conservation agriculture wheat research was pursued in Jordan, Tunisia, Algeria, Palestine and Tajikistan, through trials on-farm and on station to design, implement and demonstrate site-specific conservation agriculture packages. In Tajikistan such packages did not involve livestock whereas, in North Africa, crop-livestock integrated systems were the focus. Work in North Africa addressed alley-cropping systems, with the integration of multipurpose forage species such as *Medicago arborea, Atriplex* spp., vetch, and forage pea.

FP 5. See Sections C.2. and F.

C.2 Progress towards the achievement of outcomes

Global wheat impacts study. FP 1. To document the use of CGIAR-related and other improved wheat germplasm and its farm-level adoption, a survey was sent to 90 countries that produce at least 5,000 tons of wheat. 64 countries responded from the survey, representing 80% of the world's wheat production; responses came from 42 developing countries, covering 99% of wheat production in those regions. A full report will be published in 2015, but preliminary findings show that (1) since 1994, more than 4,600 improved wheat varieties have been released in the world and (2) there is continued and significant use of CGIAR-derived wheat germplasm in the developing world. CIMMYT-derived varieties cover 30-80% of the national wheat area in South Asian countries and, in sub-Saharan Africa, 57% of the area in Kenya and nearly 90% in Ethiopia. This study updates results of Lantican et al. (2005).

Genomic selection to speed genetic improvement in wheat breeding. FP 2. Genomic selection offers new opportunities for increasing the efficiency of plant breeding programs. In 2014, a significant body of reports in important science journals documented WHEAT scientists' substantive progress toward applying the approach.¹⁰ Topics cover included (i) genomic selection for quantitative adult plant stem rust resistance in wheat; (ii) simulations to evaluate the effectiveness of different genotyping and phenotyping strategies to enable genomic selection in early generation individuals; (iii) the results of genomic prediction in CIMMYT maize and wheat breeding programs; and (iv) use of classification algorithms for genomic-enabled prediction.

Molecular analytics to unlock new traits in massive seed collections. FP 2. As part of Seeds of Discovery (SeeD), more than 40,000 CIMMYT wheat gene bank accessions have been characterized through the DArTseq genotyping-by-sequencing (GBS) platform.¹¹ A high-density GBS consensus map developed in 2014 is an essential prerequisite for analyzing the GBS data of a large diversity panel and will facilitate

¹⁰ (i) <u>doi: 10.3835/plantgenome2014.02.0006</u>; (ii) <u>doi: 10.2135/cropsci2013.03.0195</u>; (iii) <u>doi:10.1038/hdy.2013.16</u>; (iv) <u>doi:10.1038/hdy.2013.144</u>.

¹¹ <u>http://www.diversityarrays.com/dart-application-dartseq</u>

the genetic dissection of important quantitative traits either by linkage mapping or by genome-wide association mapping.

South Asian wheat improvement with NARS partners is gaining speed (FP 3, South Asia). Twenty-one superior, resistant varieties were released in South Asia; two in Bhutan. More than 1,100 new lines qualified for inclusion in national and state trials in Bangladesh, India and Nepal. Molecular markers were identified for heat tolerance and for enhanced Zn/Fe concentrations in grain. Lines with tolerance to spot blotch were identified. Seed produced (5,639 tons) was disseminated among farmers though linkages with public and private sector institutions and also using participatory approaches.

Conservation agriculture in Ethiopia. FP 4. In June 2014, CIMMYT-Ethiopia joined the Ethiopian Highlands project of Africa RISING (see 'Africa Research in Sustainable Intensification for the Next Generation'; <u>http://africa-rising.net/where-we-work/ethiopian-highlands/</u>). Using a strong participatory research methodology, the Ethiopian Highlands project will identify technologies and management practices for the sustainable intensification of crop-livestock systems in the Ethiopian Highlands. CIMMYT is providing expertise on soil and water conservation (conservation agriculture and raised bed systems), small-scale mechanization (seeding, threshing and water pumping using two-wheel tractors), participatory variety selection (wheat, food barley, malt barley, potato and fababean) and community seed production.

Continued investment in seed systems (FP 5, Afghanistan, Ethiopia, Kenya). In **Afghanistan**, the ACIARfunded wheat improvement and seed systems project supported the production of 24,000 tons of certified seed derived from high-yielding, disease resistant CIMMYT/ ICARDA germplasm and which is enough to sow 190,000 hectares. A second phase of this project was granted also based on the donorfunded impact assessment.¹² More information about Phase 1 impact is described in Section C.3. In **Kenya**, 165 tons of seed of 10 improved varieties was supplied to 2,194 farmers; improved rust resistant varieties were demonstrated and early-generation seed multiplication done.

C.3 Progress towards impact

An overview of impact studies published 2005-2014 is shown in Annex 4, p.22.

Speeding seed production and deployment. In **Afghanistan**, seed systems research has increased supply productivity by between 65% and 79%. In **Pakistan**, through the Wheat Production Enhancement Program (<u>WPEP</u>) involving the Pakistan Ministry of Agriculture and USDA and CIMMYT, 48 tons of seed of Ug99 resistant varieties and 509 tons of seed of other rust-resistant varieties were produced and distributed to other national programs, seed suppliers and farmers. In **Ethiopia**, 165 tons of seed of 10 improved, disease-resistant bread and durum wheat varieties, along with wheat production manuals, was distributed to more than 2,000 resource-poor farmers in rust-hit areas of Arsi and Bale.

Turkey-CIMMYT-ICARDA International Winter Wheat Improvement Program (IWWIP). Four new IWWIP varieties were released in 2014 in Azerbaijan, Iran, Turkmenistan and Turkey. Varieties released by IWWIP now total 63 and are sown on 2 million hectares. CRP WHEAT is providing much needed improved wheat lines tolerant to frost and salinity for the Dryland Systems CRP Aral Sea Action Site and yellow rust resistant varieties for the Dryland Systems CRP Fergana Valley Action Site.

Yields of current vs historical varieties. In a two-year study of genetic yield potential with Windows 1&2 support comparing historical and current bread wheat and durum varieties under environments including heat and drought and covering 48 years. The 2014 release Borlaug 100 shows an 18% grain

¹² <u>http://aciar.gov.au/files/ias85-afghanistan-web-final.pdf</u>

yield advantage over Seri 82 under optimal conditions. The new Indian variety DPW621-50 showed a 14% yield advantage over PBW343 and there have been significant increases in durum wheat yields.

Sustainable intensification feasible for in South Asia. Through its research platforms, the Cereal Systems Initiative for South Asia (<u>CSISA</u>) has conclusively demonstrated sustainable intensification pathways for enhanced crop productivity (11%), increased profitability (32%), reduced energy investments (-46%) and markedly reduced irrigation requirements (-71%).¹³

D. Gender research achievements

D.1 Selected gender research achievements

Understanding gender in wheat-based livelihoods for enhanced WHEAT R4D impact in Afghanistan, Pakistan and Ethiopia, an ambitious new project accepted for funding by <u>Germany's Federal Ministry</u> for Economic Cooperation and Development in 2014, aims to contribute to reduction in rural poverty and increased food security by shaping and targeting research and development activities related to wheat, in ways that increase the empowerment of disadvantaged groups, in particular poor women and youth in wheat-based systems, and help unleash their potential. The purpose of the project is to make WHEAT research and development partners appreciate gender integration and social inclusion as an opportunity to enhance impact, and use the project's outputs to strengthen the integration of gender and social inclusion in wheat technology development, adaptation and diffusion.

Under the **Global Study on Gender Norms, Agency and Innovation** in Agriculture and Natural Resource Management (Gennovate), WHEAT completed data collection for seven case-studies in 2014, three in Morocco and four in Uzbekistan. In addition, under the WHEAT Competitive Partner Grant scheme, a grant to undertake a comprehensive diagnosis of gender relations in key wheat target regions of South Asia and development of guidelines for gender-responsive wheat-based systems development was established. Implementation of the project is led by Glasgow Caledonian University in collaboration with local partners in Bangladesh, India, Nepal and Pakistan, and involves the organization and operationalization of 30 case studies under Gennovate.

D.2 Success and challenges in mainstreaming gender research

A revised version of the <u>WHEAT gender strategy</u> was endorsed by the WHEAT management committee in December. Other successes related to gender mainstreaming include increased demand from scientists and research teams for gender inputs for wheat research-for-development (R4D), as well as increased efforts to integrate gender considerations into technical wheat R4D. Numerous challenges to mainstream gender in WHEAT remain, in particular related to bilateral funding and insecurity in W1&2 funding, which significantly limits the capacity to increase CRP gender staffing for mainstreaming efforts.

Gender performance self-assessment as per Annex 2, p.20: Approaching requirements.

E. Partnership building achievements

E. 1. Selected partnership-building achievements

Global Partnership Meeting. In a two-day meeting in Istanbul, Turkey, in December 2014, more than 100 participants from various sectors and institutions around the globe helped shape the beginnings of a proposal for WHEAT's second phase (2017-21), including the scope and outcomes of Flagship Projects, partnerships needed, and next steps for project designs and leadership.

¹³ See p. 8 of <u>this CSISA report</u>.

Boosting wheat production in Africa to stem costly imports. Through the <u>SARD-SC</u> wheat project, during 2014 ICARDA operationalized 3 hubs and 15 innovation platforms in Ethiopia, Nigeria and Sudan. The impressive performance of new heat-tolerant wheat varieties in Nigeria and Sudan with yields of 4-6 t/ha– significantly more than 1-2 t/ha average of traditional varieties – generated a key policy shift in both countries. Accordingly, wheat has been included as a priority in the Nigerian Government's Agricultural Transformation Agenda (ATA) and domestic production is targeted as a solution for curbing ever growing import dependence and for ensuring food security. The Government of Nigeria has launched a national target to scale-up best practices, expand wheat area from the existing 70,000 ha to 340,000 ha, and reduce the wheat importation by 45% in the coming five years. Likewise, based on the high yield realized by wheat farmers participating in the six project IP sites in Sudan, the Sudan Government formally announced intentions to scale up the IP approach through the national extension program, expanding the country's wheat area from the present 137,000 ha to 340,000 ha in 2015/16 and boosting domestic production to reduce costly imports.

Bundling wheat improvement resources in Central Asia. The Kazakhstan-Siberian Network on Wheat Improvement (KASIB) established and coordinated by CIMMYT currently unites 19 breeding programs in Eurasia (Kazakhstan, Western Siberia, Ural, Altai and Volga), covering more than 20 million ha of wheat. KASIB activities over 14 years have generated more than 15,000 wheat lines and tested 469 advanced spring bread wheat varieties and 136 advanced spring durum wheat varieties. More than 10 varieties are presently submitted to State Testing Committees.

Partnership for climate change modeling. Successful collaboration with the wheat modeling group of the Agricultural Model Intercomparison and Improvement Project (AgMIP--Senthold Asseng, University of Florida, USA) has (1) assessed the state of the science on modeling climate change impacts for wheat, (2) generated constructive methods and novel algorithms for representing wheat's response to high temperatures and other stresses, and (3) leveraged a relatively small budget to conduct global research documented in a high-impact journal (see *Nature Climate Change* article cited on p.3).

E.2 Strategic partnerships

IWYP(International Wheat Yield Partnership) Launched (USAID funded). A key WHEAT contribution to this BBSRC/USAID-led project to increase genetic potential by 50% by 2030 is MEXPLAT (more detail about that platform in section C1).

JIRCAS-funded FP 2 - Genetically modified (GM) wheat field trial in Mexico. Evaluation of 5 genepromoter combinations for drought tolerance in wheat and rice has been conducted using an integrated, multidisciplinary approach spanning plant molecular biology, physiology, and breeding collaborations JIRCAS- RIKEN, CIAT, IRRI and CIMMYT. (To read CIMMYT's policy statement on genetically modified crops, <u>click here</u> and scroll down.)

E.3 Interactions with other CRPs

Agriculture Innovation in Pakistan: Multi-Partner, including multi-CRP. Under the Agricultural Innovation Program (AIP) for Pakistan, WHEAT interacts with the A4NH, MAIZE and Livestock and Fish CRPs on feed quality of wheat and maize varieties. AIP also works with WHEAT, MAIZE and GRiSP on rice-wheat-based systems; with the Policies, Institutions and Markets to establish research boards in Pakistan; and with Dryland Systems and ICARDA in wheat agronomy. Through AIP also works through competitive grants with Legumes. AIP partners with the World Vegetable Center (AVRDC) on vegetables and with UC Davis in research on fruit crops. Supported by CIMMYT efforts, ILRI, IRRI, AVRDC, and UC Davis opened offices in Pakistan and CIAT is signing an MoA with CIMMYT. Finally, AIP has created a <u>new innovation platform</u> with Pakistan's National Rural Support Program (NRSP) for wheat varietal

evaluation, popularization and deployment, on-farm agronomic interventions and community-based seed production enterprises, thereby opening a key avenue to reach large numbers of smallholders, tenants, women headed households and vulnerable people in food-deficit areas not served by mainstream public or private sector organizations.

HarvestPlus (A4NH) and WHEAT: A two-way street. Research under FP3 is making use of HarvestPlus outputs directly and indirectly. Breeding for enhanced micronutrients involves landraces and other wild ancestors. Selected novel alleles identified for traits such as yield and stress tolerance are mobilized into WHEAT breeding programs. Breeding for high Zn and for yield is handled as in the main breeding program up to the F_5 stage. Thus have HarvestPlus cross derivatives entered the Elite Selection Wheat Yield Trials (see also 'Nutritious wheat', p.5).

F. Capacity building

Gender equality on the rise in training. In 2014, with funding from WHEAT, **CIMMYT Global Wheat Program (GWP)** training courses involved 89 junior and mid-career scientists from all over the world; nearly 50% were women. The courses ranged from the basic wheat improvement course in Mexico (15 countries represented) to the regional statistics course on biotechnology and molecular breeding in Tunisia (6 countries represented) with ICARDA and IRESA (the Ministry of Agriculture) Wheat training courses have encouraged female participation by adjusting courses and accommodating for those who are unable to stay for extended periods. **The Jeanie Borlaug Award Program for Women in Triticum** (WIT) was established by GWP and Cornell University as part of the Durable Rust Resistance in Wheat Project (<u>DRRW</u>). **ICARDA** hosted 279 trainees in 2014; 14% female. Courses ranged from quality seed production to molecular breeding, crop improvement and supplemental irrigation and were held in Sudan, Nigeria, Morocco, Ethiopia, Tunisia and Iraq. In addition to group courses, ICARDA held on-site farm workshops and study missions for students. A total of 13 newly enrolled PhD and MSc students participated in courses in 2014; 77% were female.

First phase of the Arab Food Security Project (2011-14). 25,700 farmers have benefited from field days, farmer schools and travelling workshops, with an average 28% increase in wheat yield across all countries from large-scale on-farm demonstrations of improved wheat varieties and agronomic practices. Mechanized raised-bed planting for smallholders provided an average 25% savings in irrigation water, a 30% increase in wheat yield and a 74% improvement in water-use efficiency on farmers' fields in Egypt. Substantial improvements in rainfed systems included a 45% gain in yields in Yemen and gains of 24% in Jordan and Tunisia. No-till systems increased wheat yields 16% in Syria, 20% in Jordan and 50% in Morocco. The project's Young Scientist Training Program mentored and trained 34 young scientists. Work was funded by the Arab Fund for Economic and Social Development, Kuwait Fund for Arab Economic Development, Islamic Development Bank and OPEC Fund for International Development, and is now set to scale out the successes. Full story here.

G. Risk management

In 2014, a risk management matrix was created to regularly assess and manage risks related to the delivery of results by the CRP. This matrix identifies a number of risks related to asset management, compliance, general management, change management, finance and technology. Based on the risk assessment, specific mitigation measures were identified to manage these risks.

The three major risks for the CRP are: (1) delayed transfer of W1&2 funds, which directly affects CRP research and development operations; (2) non-fulfilled obligations by the partners for commissioned and competitive grants; (3) lack of a systematic and integrated approach for monitoring and evaluation at the outcome level.

To mitigate risk (1), the WHEAT-Management Committee agreed to prioritize CGIAR-led research over partner and management budgets, and to maintain the WHEAT partner budget as the most flexible component of the budget. WHEAT continues to sign only one-year partner grant contracts, to manage partner expectations and minimize any delays of payments to them. As for risk (2), WHEAT has hired a Senior Monitoring, Evaluation and Learning Specialist, shared with the MAIZE, to strengthen the CRP monitoring and evaluation system. Other CIMMYT initiatives also contribute to minimizing this risk, including the implementation of a documentation and monitoring system.

H. Lessons learned (including monitoring CRP progress)

Notwithstanding the positive outcomes achieved, additional efforts are needed to increase the quality of project management and of monitoring and evaluation at both project and program levels.

To respond to this need, both CIMMYT-led CRPS improved the standardized process and associated tools for project planning and design, including clear roles and responsibilities for each step and decision. It has also identified key steps where specialists (e.g., gender, monitoring and evaluation, communications) should be involved to provide advice. At the institutional level, CIMMYT has launched a new component of its Research Management System to support project planning and to record key documentation and decisions.

The CRP foresees a number of future efforts to support this need, including building capacity in the development of theories of change, impact pathways and monitoring and evaluation plans, and the development of a monitoring, evaluation and learning strategy.

H.1 Level of confidence of the response to the key performance indicators

The information reported in Table 1 is obtained from detailed data found in a variety of sources, including project technical reports and institutional databases. WHEAT is confident in the quality of the indicator information supplied. However, the program will continue to improve the systematic approach to collecting the quantitative evidence and other types of performance or progress data, across the WHEAT project portfolio, to improve the process and time required to collect and analyze the information.

WHEAT has reorganized its strategy around five Flagship Projects in 2014 encompassing the Strategic Initiatives of the original proposal following a standardization of CRP structure. These changes have been reflected in Table 1, under indicators 1, 2 and 3.

The program also endorses other CRPs' positions on the importance of clarifying the definitions of some of the SLO- and IDO-level indicators to support consistent and reliable reporting across CRPs.

H.2 Unintended results and innovative initiatives

There were no unintended results.

Part II: Financial report

Figure 1. Total WHEAT budget vs. expenditures, 2014 (US \$ millions): L111 and L136



Figure 2. POWB approved by Flagship Project, 2014.





Figure 3. Actual expenditures by Flagship Project, 2014.

Figure 4. Gender Summary by Flagship Project, 2014 (US \$ millions).



Table 1. Annual Funding Summary (Totals for Windows 1 and 2 combined): L106

		Windows 1&2	Window 3	Bilateral Funding	Total Funding
	CIMMYT :				
1	CGIAR	15,340	-		15,340
2	ACIAR	· · · · ·	230	-	230
3	Arcadia		-	648	648
4	Barcel, S.A. de C.V.		-	50	50
5	BMGF	-	2,103	-	2,103
6	Cornell	-	-	2,905	2,905
7	GIZ	-	-	346	346
8	ICAR, India	-	403	-	403
9	ICARDA	-	-	96	96
10	Japan	-	160	69	228
11	Kansas	-	-	317	317
12	OFID	-	-	54	54
13	SAGARPA	-	-	2,805	2,805
14	Turkey	-	226	-	226
15	UOT	-	-	58	58
16	USAID	-	3,456	-	3,456
17	Other <50k	-	70	79	149
	ICARDA:				
18	International Maize and Wheat Improvement	2,854		291	3,145
19	Germany			386	386
20	Kuwait Fund			498	498
21	Swedish University of Agricultural Sciences (via the Swedish Researc	h Council)	79	79
22	African Development Bank (AfDB) through IIT.	Α		2,027	2,027
23	Agricultural Research Center- Egypt (ARC)			50	50
24	Arab Fund for Economic and Social Developm	ent (AFESD)		752	752
25	Cornell University			501	501
26	European Commission		872		872
27	Grains Research & Development Corporation	(GRDC)		281	281
28	Islamic Development Bank (ISDB)			98	98
29	Japan		66		66
30	NSW Department of Primary Industries (DPI)			80	80
31	The Japan International Cooperation Agency (JICA)		204	204
32	Turkey		239		239
33	United States Agency for International Develo	opment (USAID)	1,607		1,607
34	United States Department of Agriculture (USD	A)		810	810
35	Other <50k			87	87
Total for CRP 3.1: WHEAT		18,194	9,431	13,571	41,196

	Windows 1&2	Window 3	Bilateral Funding	Center Funds	Total Funding	Windows 1&2	Window 3	Bilateral Funding	Center Funds	Total Funding
Total CRP 3.1: WHEAT		P	OWB Approved Budg	et				Actual		
Personnel	5,221	2,853	6,566	-	14,640	5,558	1,969	3,583	-	11,110
Collaborators Costs - CGIAR Centers	3,044	2,789) -	-	5,833	4,497	1,567	-	-	6,064
Collaborator Costs - Partners	178	2,210) 1,372	-	3,760	256	1,498	1,812	-	3,566
Supplies and services	7,701	3,472	2 5,289	-	16,462	3,006	2,408	5,345	-	10,759
Operational Travel	579	399) 1,256	-	2,234	606	356	815	-	1,777
Depreciation	1,019	396	3 1,168	-	2,583	2,118	511	635	-	3,265
Sub-total of Direct Costs	17,742	12,119	15,651	-	45,511	16,042	8,309	12,189	-	36,540
Indirect Costs	1,802	1,585	5 1,807	-	5,194	2,152	1,123	1,381	-	4,656
Total - All Costs	19,544	13,704	17,457	-	50,705	18,194	9,431	13,571	-	41,196
LESS Coll Costs CGIAR Centers	(3,044)	(2,789)) -	-	(5,833)	(4,497)	(1,567)	-	-	(6,064)
Total Net Costs	16,500	10,915	5 17,457	-	44,872	13,697	7,865	13,571	-	35,133

Table 2. Annual financial summary by natural classification (US \$ 000s): L121

Windows 1 & 2

Bilateral Funding Center Funds Total Funding

Total CRP 3.1: WHEAT	Unspent/Variance										
Personnel	(338)	884	2,983	-	3,529						
Collaborators Costs - CGIAR Centers	(1,453)	1,223	-	-	(231)						
Collaborator Costs - Partners	(78)	712	(440)	-	194						
Supplies and services	4,695	1,064	(56)	-	5,703						
Operational Travel	(27)	43	441	-	458						
Depreciation	(1,100)	(115)	533	-	(682)						
Sub-total of Direct Costs	1,699	3,810	3,462	-	8,971						
Indirect Costs	(350)	463	425	-	538						
Total - All Costs	1,349	4,273	3,887	-	9,509						
LESS Coll Costs CGIAR Centers	1,453	(1,223)	-	-	231						
Total Net Costs	2,803	3,050	3,887	-	9,740						

Window 3

Table 3. WHEAT partnership report, 2014 (US \$ 000s): L211

TOTAL	FOR CRP "X.X"				Actual Ex	penses - This Year	Science for a food secure future		
	Institute Acronym	Institute Name	<u>Country</u>	Windows 1 & 2	Window 3	Bilateral	Center Funds	TOTAL	
	CIMMYT :								
1	ASARECA	Association for str	eng UGANDA	100	-	-	-	100	
2	BARI	Bangladesh Agricu	Itur BANGLADESH	-	238	(2)	-	236	
3	BIU	BANARAS HINDU U	NIVIINDIA	184	14	-	-	198	
4	CAS	CREATIVE AGRI SOL	UTI(CHINA	94	-	-	-	94	
5	CU	CORNELL UNIVERSI	TY USA	-	-	74	-	74	
6	DATECH	DIVERSITY ARRAYS	TECIAUSTRALIA	90	-	25	-	115	
7	DWR	INIDIAN INSTITUTE	OF INDIA	118	-	-	-	118	
8	ICAR	Indian Council of	Agric INDIA	-	39	235	-	274	
9	ICARDA	INTERNATIONAL CE	NTE ARAB REPUBLIC	986	-	-	-	986	
10	IDE	INTERNATIONAL DE	EVEL BANGLADESH	-	2/1	-	-	2/1	
11	IFPRI	INTERNATIONAL FO		-	80	-	-	80	
12	IISERK			205	- 01	-	-	203	
10				-	1/1	-	-	1/1	
14				- 672	141	-	-	141 672	
16	INIFAD	INSTITUTO NACION		0/5		168		168	
17	IRRI	INTERNATIONAL RI			1 227			1.227	
18	IWWIP	INTERNATIONAL W	INTETURKEY	36	69	-	-	105	
19	KRIGBSPK	KASHKADARYA RES	EAR(UZBEKISTAN	120	-	-	-	120	
20	LAU	LANCASTER UNIVER	RSITYUK	466	-	-	-	466	
21	RRL	ROTHAMSTED RESE	ARC UK	374	-	-	-	374	
22	RUM	REGENTS OF THE U	NIVEUSA	200	104	-	-	304	
23	TARI	TIGRAY AGRICULTU	IRAL ETHIOPIA	-	-	53	-	53	
24	TG	TRAIT GENETICS Gr	nbH GERMANY	187	-	-	-	187	
25	UACHL	UNIVERSIDAD AUS	TRAL CHILE	-	-	55	-	55	
26	UCGC	THE UNIVERSITY CO	DURTUK	473	-	-	-	473	
27	WAUR	WAGENINGEN UR	NETHERLANDS	307	-	-	-	307	
28	Others les than 50k	:		67	178	96	-	341	
	ICARDA:							-	
29	EIAR	Ethiopian Institute	e of Ethiopia		335	41		376	
30	CIMMYT	The International	Maize and Wheat Improve	ement Center		300		300	
31	University of Bonn	University of Bonn	Gemany			283		283	
32	ARC	Agricultural Resea	irch i Sudan		13	79		92	
33	INRA	Institut National o	le la Morocco		53	33		87	
34	ARC	Agricultural Resea	irch i Egypt		21	60		81	
35	IWWIP	International Win	ter \ Turkey		71			71	
36	NIBGE	National Institute	of E Pakistan			56		56	
37	Others les than 50k			15	128	255		398	
								-	
								-	
То	otal for CRP			4,753	3,064	1,811	-	9,629	

Annex 1. Indicator of progress in 2014.

CRP all-FP completion rate	97.1%	
CRP all-FP reporting		
completion	91.3%	

	Flagship 1		1	Flagship 2			Flagship 3			Flagship 4			Flagship 5			Flagship 2&3				
	CoA 1.1	CoA 1.2	CoA 1.3	CoA 2.1	CoA 2.2	CoA 2.3	CoA 2.4	CoA 3.1	CoA 3.2	CoA 3.3	CoA 3.4	CoA 3.5	CoA 4.1	CoA 4.2	CoA 4.3	CoA 5.1	CoA 5.2	CoA 5.3	CoA 2./3.6 (acc. 50/50)	CoA 2./3.7 (acc. 50/50)
Completion per CoA	92.5%	83.8%	100%	100%	100%	97.2%	91.8%	95.8%	96.4%	100%	91.9%	99.5%	100%	100%	100%	99.8%	100%	100%	91.4%	100%
Completion		1																		
per FP	92.1% 96.9%		.9%				96.6%				100.0%			99.9%		95.	7%			
Reporting																				
completion		84.6%			90).6%				89.7%			97.4%			94.1%		90.0%		

CRP indicators of progress, with glossary and targets.

CRPs			Deviation narrative (if	20	13	2	014	20	15
concerned by this indicator	Indicators	Glossary & Comments	actual is more than 10% away from target)	Target	Actual	Target	TOTAL	Target	Actual
All	1. Number of flagship "products" produced by CRP	Glossary: These are frameworks and concepts they should be likely to change the way stakeholders along the impact pathway allocate resources and/or implement activities change the way these stakeholders think and act. For the CRP MAIZE, each Flagship Project is a flagship "product.	Following a standardization of CRP structures, the MAIZE strategy was reorganized around five Flagship Projects (FPs) in 2014, encompassing the nine Strategic Initiatives of the original MAIZE proposal	10	10	5	5	5	
All	2. % of flagship products produced that have explicit target of women farmers/NRM managers	Included in FPs: FP1 - Maximizing value for money, social inclusivity through prioritizing WHEAT R4D FP3 - Partnership to accelerate genetic gain in farmers field FP4 - Sustainable intensification FP5 - Human and institutional capacities, scaling out and up	Following a standardization of CRP structures, the MAIZE strategy was reorganized around five Flagship Projects (FPs) in 2014, encompassing the nine Strategic Initiatives of the original MAIZE proposal	3	4	4	4	4	
All	3. % of flagship products produced that have been assessed for likely gender-disaggregated impact	Included in FPs: FP1-Maximizing value for money, social inclusivity through prioritizing WHEAT R4D FP3 - Partnership to accelerate genetic gain in farmers field FP4 - Sustainable intensification FP5 - Human and institutional capacities, scaling out and up	Following a standardization of CRP structures, the MAIZE strategy was reorganized around five Flagship Projects (FPs) in 2014, encompassing the nine Strategic Initiatives of the original MAIZE proposal	10	10 of which 2 more in- depth	4	4	4	
All	4. Number of "tools" produced by CRP	Glossary: These are significant decision-support tools, guidelines, training manuals, software, and/or videos that are significant in that they should be likely to change the way stakeholders along the impact pathway allocate resources and/or implement activities		522 changed to 25	23 (17 co- developed with other CRPs)	26	39	25	
All	 % of tools with explicit target of women farmers 	Tools target men and women users equally							
All	6. % of tools assessed for likely gender-disaggregated impact	Tools are not assessed individually but at flagship product level							
All	7. Number of open access databases maintained by CRP	IWIS, GRIS (wheatpedigree.net), IWIP.org, rusttracker.org, Wheat Atlas and Wheat Doctor (taken off the web Dec 2013) Not included : Cereal Knowledge Bank, http://www.knowledgebank.irri.org/wheat.ht, maintained by IRPI (with 118,671 unique pageviews)		5	6	Tobe reviewed	11	5	
All	 Total number of users of these open access databases 	Unique visitors (not included : Cereal Knowledge Bank)			124,450	54,767	119,832	125,000	
All	9. Number of publications in ISI journals produced by CRP			121	121 (23 jointly with other CRPs)		107	120	
1,2,3, 4, 6	10. Number of strategic value chains analyzed by CRP	Ethiopia: addition 2013; 3 for India/CSISA (wheat value chains in states Bihar, Haryana and Madhya Pradesh)		5	4	2	10	5	
1,5,6,7	11. Number of targeted agro- ecosystems analysed/characterised by CRP								
1,5,6,7	12. Estimated population of above-								

CRP indicators of progress, with glossary and targets (cont'd).

		WHEAT							
CRPs			Deviation narrative (if	20	13	2	014	20	15
concerned by this indicator	Indicators	Glossary & Comments	actual is more than 10% away from target)	Target	Actual	Target	TOTAL	Target	Actual
		TECHNOLOGIES/PRACTICES IN VARIOUS STA	GES OF DEVELOPMENT						
All	18. Number of technologies/NRM practices under research in the CRP (Phase I)	Germplasm: 184,633 Agronomy: 372 From KPI database		260,000	124,517		200,005	120,000	
All	 % of technologies under research that have an explicit target of women farmers 								
All	20. % of technologies under research that have been assessed for likely gender-disaggregated								
1,5,6,7	21Number of agro-ecosystems for which CRP has identified feasible approaches for improving ecosystem services and for establishing positive incentives for farmers to improve ecosystem functions as per CRP's recommends								
1,5,6,7	22. Number of people who will potentially benefit from plans, once finalised, for the scaling up of strategies								
All, except 2	23. Number of technologies /NRM	Germplasm: 1325 Agronomy: 76 From KPI database		2,600	2600		2,151	1,000	
1,5,6,7	24. Number of agro-ecosystems for which innovations (technologies, policies, practices, integrative approaches) and options for improvement at system level have been developed and are being field tested (Phase II)								
1,5,6,7	25. % of above innovations/approaches/ options targeted at decreasing inequality between men and women								
1,5,6,7	26. Number of published research outputs from CRP utilised in targeted agro-ecosystems								
All, except 2	27.Number of technologies/NRM practices released by public and private sector partners globally (phase III)	Agronomy: 14 From KPI database		50	46	50	24	50	

CRP indicators of progress, with glossary and targets (cont'd).

WHEAT												
CRDs			Deviation narrative (if	2013		2014		2015				
concerned by this indicator	Indicators	Glossary & Comments	actual is more than 10% away from target)	Target	Actual	Target	TOTAL	Target	Actual			
CAPACITY ENHANCEMENT AND INNOVATION PLATFORMS												
All	13. Number of trainees in short-term programs facilitated by CRP (male)	From CIMMYT Training database, plus Competitive Partner Grants and ICARDA Short-term = < 90 days		16,415	14,232 (171 with other CRPs)	17,000	4,088	3,000				
All	14. Number of trainees in short-term programs facilitated by CRP (female)	From CIMMYT Training database, plus Competitive Partner Grants and ICARDA Short-term = < 90 days		5,226	3,068 (73 with other CRPs)	To be reviewed	1,069	1,000				
All	15. Number of trainees in long-term programs facilitated by CRP (male)	From CIMMYT Training database, plus Competitive Partner Grants and ICARDA Long-term = > 90 days		40	121 (7 with other CRPs)	Tobe reviewed	44	24				
All	16. Number of trainees in long-term programs facilitated by CRP (female)	From CIMMYT Training database, plus Competitive Partner Grants and ICARDA		23	65 (4 with other CRPs)	To be reviewed	36	24				
1,5,6,7	17. Number of multi-stakeholder R4D innovation platforms established for the targeted agro-ecosystems by the CRPs			43	45	50	50	45				
POLICIES IN VARIOUS STAGES OF DEVELOPMENT												
All	28. Numbers of Policies/ Regulations/ Administrative Procedures Analyzed (Stage 1)			2	6	2	8	2				
All	 Number of policies / regulations / administrative procedures drafted and presented for public/stakeholder consultation (Stage 2) 			1			3	1				
All	30. Number of policies / regulations / administrative procedures presented for legislation (Stage 3)			o	1	1	1	1				
All	31. Number of policies / regulations / administrative procedures prepared passed/approved (Stage 4)						1	1				
All	32. Number of policies / regulations / administrative procedures passed for which implementation has begun (Stage 5)						0	1				
OUTCOMES ON THE GROUND												
All	33. Number of hectares under improved technologies or management practices as a result of CRP research	Based on impact studies for select geographies: RWC-CSISA/Laser Land Levelling (2014 study): 1.5M ha, 1.2M farmers (1988-2013); Ethiopia 90% national wheat area covered w/ CGIAR-derived varieties (2014 study, 2015 global impact study): 1.53M ha, 3.06M farmers; China impact study (2015): CGIAR-derived varieties sown 1.815M ha, 6.05M farmers (1982-2012)		1,650,000	1,650,000		4,845,000	1,815,000				
All	34. Number of farmers and others who have applied new technologies or management practices as a result of CRP research	see above		2,802,000	2,802,000		10,310,000	3,082,200				

Annex 2. Performance indicators for gender mainstreaming with targets defined.

Performance Indicator	CRP performance approaches requirements	CRP performance meets requirements	CRP performance exceeds requirements
1. Gender inequality targets defined	Sex-disaggregated social data is being collected and used to diagnose important gender-related constraints in at least one of the CRP's main target populations	Sex-disaggregated social data collected and used to diagnose important gender-related constraints in at least one of the CRP's main target populations And The CRP has defined and collected baseline data on the main dimensions of gender inequality in the CRP's main target populations relevant to its expected outcomes (IDOs)	Sex-disaggregated social data collected and used to diagnose important gender-related constraints in at least one of the CRP's main target populations And The CRP has defined and collected baseline data on the main dimensions of gender inequality in the CRP's main target populations relevant to its expected outcomes (IDOs) And CRP targets changes in levels of gender inequality to which the CRP is or plans to contribute, with related numbers of men and women beneficiaries in main target populations
2. Institutional architecture for integration of gender is in place	 CRP scientists and managers with responsibility for gender in the CRP's outputs are appointed, have written TORS. Procedures defined to report use of available diagnostic or baseline knowledge on gender routinely for assessment of the gender equality implications of the CRP's flagship research products as per the Gender Strategy CRP M&E system has protocol for tracking progress on integration of gender in research 	 - CRP scientists and managers with responsibility for gender in the CRP's outputs are appointed, have written TORS and funds allocated to support their interaction. - Procedures defined to report use of available diagnostic or baseline knowledge on gender routinely for assessment of the gender equality implications of the CRP's flagship research products as per the Gender Strategy -CRP M&E system has protocol for tracking progress on integration of gender in research And A CRP plan approved for capacity development in gender analysis 	CRP scientists and managers with responsibility for gender in the CRP's outputs are appointed, have written TORS and funds allocated to support their interaction. - Procedures defined to report use of available diagnostic or baseline knowledge on gender routinely for assessment of the gender equality implications of the CRP's flagship research products as per the Gender Strategy -CRP M&E system has protocol for tracking progress on integration of gender in research And A CRP plan approved for capacity development in gender analysis And The CRP uses feedback provided by its M&E system to improve its integration of gender into research

Annex 3. How WHEAT delivers towards impacts (CGIAR System Level Outcomes).



Figure A3-1. Overview of FP outputs delivering towards sub-IDOs and SLOs.

Figure A3-2. Wheat CRP impact pathways.



Annex 4. WHEAT-related impact studies published during 2004-2014.

A 2014 study identified total of 105 relevant reports for 2004-14, including 74 with a focus on CRP-Wheat and 31 studies across-CRPs. Most directly involved CIMMYT and/or ICARDA scientists as (co-) authors. A few were fully external, published and available in the public domain, examples including ACIAR's commissioned external impact assessment of CIMMYT implemented projects in Afghanistan (Jilani et al., 2013) and an independent journal paper evaluating CIMMYT achievements in NW Mexico – the cradle of the green revolution (Nalley et al., 2010). There are also hybrid publications – published by a CGIAR center but led by external scholars (e.g. Gollin, 2006).

Most relate to adoption and impacts of technological interventions associated with CIMMYT, ICARDA and/or CRP-Wheat for farmers and, to a more limited extent, consumers. The 105 studies amount to an average of 9.5 publications per year. About half were peer reviewed and a third (32%, 3.1 p.a.) were published in recognized journals (Table 1). Contrasting the pre-CRP years (2004-11) to the CRP-years (2012-14), the total number of annual publications increased from 9.1 to 10.7 p.a (17%). The number of peer reviewed publications doubled from 3.5 to 7.0 p.a., representing respectively 38% and 66% of the period's publications, although the number of recognized journals increased more modestly. For fuller details of this report and the publications listed, <u>click here</u>.

Annex 5. 2014 WHEAT Publications

Articles in refereed journals

- Akhmetova, A., Zelenskiy, Y., Karabayev, M., Morgounov, A.I. (2014). The yield and resistance to rust of KASIB nurseries, Accepted for print.
- Al-Saleh, A., Baloch F. S., Derya M., Azrak M., Kilian B., Özkan H. and Nachit M. (2014). Genetic Linkage
 Map of Anatolian Durum Wheat Derived from a Cross of Kunduru-1149 × Cham1. Plant Mol Biol Rep.
 DOI 10.1007/s11105- 014-0749-6.
- Aranjuelo, I., Molero, G., Avice, J.C., Bourguignon, J.(2014). A novel method for determination of the 15N isotopic composition of Rubisco in wheat plants exposed to elevated atmospheric carbon dioxide, Physiologia Plantarum, In press. http://onlinelibrary.wiley.com/doi/10.1111/ppl.12294/abstract
- Awais Rasheed, A., Xianchun Xia, Ogbonnaya, F., Mahmood, T., Zongwen Zhang, Mujeeb-Kazi, A., He Zhonghu (2014). Genome-wide association for grain morphology in synthetic hexaploid wheats using digital imaging analysis, BMC Plant Biology, 14, 128, 21 p. <u>http://www.biomedcentral.com/1471-2229/14/128</u>
- Babu, R., Palacios Rojas, N., Prasanna, B.M. (2014). Biofortified maize a genetic avenue for nutritional security, Translational genomics for crop breeding: Abiotic stress, yield and quality. Vol. 2, p. 161-176. <u>http://onlinelibrary.wiley.com/doi/10.1002/9781118728482.ch10/summary</u>
- Bari, A., Amri A., Street K., Mackay M., De Pauw E., Sanders R., Nazari K., Humeid B., Konopka J. and Alo F. (2014). Predicting resistance to stripe (yellow) rust (Puccinia striiformis) in wheat genetic resources using focused identification of germplasm strategy. The Journal of Agricultural Science, available on CJO2014. DOI: 10.1017/S0021859613000543.
- Basnet, B.R., Singh, R.P., Ibrahim, A.M.H., Herrera-Foessel, S.A., Huerta-Espino, J., Caixia Lan, Rudd, J.C. (2014). Characterization of Yr54 and other genes associated with adult plant resistance to yellow rust and leaf rust in common wheat Quaiu 3, Molecular Breeding, 33, 2, 385-399.
 http://link.springer.com/article/10.1007/s11032-013-9957-2
- Baudron, F., Delmotte, S., Corbeels, M., Herrera, J.M., Tittonell, P. (2014). Multi-scale trade-off analysis of cereal residue use for livestock feeding vs. soil mulching in the Mid-Zambezi Valley, Zimbabwe, Agricultural Systems, In press. http://www.sciencedirect.com/science/article/pii/S0308521X14000328
- Bellarby, J., Stirling, C., Vetter, S.H., Kassie, M., Kanampiu, F., Sonder, K., Smith, P., Hillier, J. (2014).
 Identifying secure and low carbon food production practices: A case study in Kenya and Ethiopia, Agriculture, Ecosystems and Environment, 197, 137-146.
 http://www.sciencedirect.com/science/article/pii/S0167880914003752
- Bishaw, Z., Struik P.C. and van Gastel A.J. G. (2014). Assessment of on-farm diversity of wheat varieties: Evidence from farmer's fields in Ethiopia. African Journal of Agricultural Research 9(39): 2948-2963.
- Caixia Lan, Singh, R.P., Calvo-Salazar, V., Herrera-Foessel, S.A. (2014). Genetic analysis of resistance to leaf rust and stripe rust in wheat cultivar Francolin#1, 2014, Plant Disease, 98, 9, 1227-1234. http://apsjournals.apsnet.org/doi/abs/10.1094/PDIS-07-13-0707-RE

- Chand, R., Kumar, M., Kushwaha, C., Shah, K., Joshi, A.K. (2014). Role of melanin in release of extracellular enzymes and selection of aggressive isolates of bipolaris sorokiniana in barley, Current Microbiology, 69, 2, 202-211. <u>http://link.springer.com/article/10.1007%2Fs00284-014-0559-y</u>
- Chhokar, R.S., Sharma, R.K., Gathala, M.K., Pundir, A.K. (2014). Effects of crop establishment techniques on weeds and rice yield, Crop Protection, 64, 7-12. http://www.sciencedirect.com/science/article/pii/S0261219414001793
- Chung, U., Gbegbelegbe, S.D., Shiferaw, B., Robertson, R., Jin I. Yun, Kindie Tesfaye Fantaye, Hoogenboom, G., Sonder, K. (2014). Modeling the effect of a heat wave on maize production in the USA and its implications on food security in the developing world, Weather and Climate Extremes, 6-May, 67-77. <u>http://www.sciencedirect.com/science/article/pii/S2212094714000668</u>
- Crespo-Herrera, L.A., Akhunov, E., Garkava-Gustavsson, L., Jordan, K.W., Smith, C.M., Singh, R.P., Ahman, I. (2014). Mapping resistance to the bird cherry-oat aphid and the greenbug in wheat using sequence-based genotyping, Theoretical and Applied Genetics, 127, 9, 1963-1973. <u>http://link.springer.com/article/10.1007%2Fs00122-014-2352-5</u>
- Crossa, J., Perez, P., Hickey, J.M., Burgueño, J., Ornella, L., Ceron-Rojas, J., Zhang, X., Dreisigacker, S., Babu, R., Yongle Li, Bonnett, D.G., Mathews, K.L. (2014). Genomic prediction in CIMMYT maize and wheat breeding programs, Heredity, 112, 48-60. http://www.nature.com/hdy/journal/vaop/ncurrent/abs/hdy201316a.html
- Crossa, J., Vargas, M., Cossani, C.M., Alvarado, G., Burgueño, J., Mathews, K.L., Reynolds, M.P. (2014). Evaluation and interpretation of interactions, Agronomy Journal, 105, 1, 1-12. <u>https://www.soils.org/publications/aj/articles/0/0/agronj2012.0491</u>
- Dababat, A.A. (2014). Biological and chemical dependent systemic resistance and their significance for the control of root-knot nematodes, Nematology, 16, 8, 917-927. http://booksandjournals.brillonline.com/content/journals/10.1163/15685411-00002818
- Dababat, A.A., Erginbas-Orakci, G., Toktay, H., Imren, M., Akin, B., Braun, H-J, Dreisigacker, S., Elekcioglu, I.H., Morgounov, A.I. (2014). Resistance of winter wheat to Heterodera filipjevi in Turkey, Turkish Journal of Agriculture and Forestry, 38, 180-186.
- Dababat, A.A., Imren, M., Erginbas-Orakci, G., Ashrafi, S., Yavuzaslanoglu, E., Toktay, H., Pariyar, S.R., Elekcioglu, I.H., Morgounov, A.I., Mekete, T. (2014). The importance and management strategies of cereal cyst nematodes, Heterodera spp., in Turkey, Euphytica, In press. <u>http://link.springer.com/article/10.1007/s10681-014-1269-z</u>
- Dababat, A.A., Orakci, G.E., Toktay, H., Imren, M., Akin, B., Braun, H.J., Dreisigacker, S., Elekcioglu, I.H., Morgounov, A.I. (2014). Resistance of winter wheat to Heterodera filipjevi in Turkey, Turkish Journal of Agriculture and Forestry, 38, 180-186. <u>http://journals.tubitak.gov.tr/agriculture/issues/tar-14-38-</u> 2/tar-38-2-3-1305-47.pdf
- Dababat, A.A., Pariyar, S.R., Nicol, J.M., Erginbas-Orakci, G., Goll, M., Watrin, C., Duveiller, E., Braun, H.J., Cabrera, J.A., Sikora, R.A. (2014). Influence of thiabendazole seed treatment on the integrated control of Heterodera filipjevi on six wheat genotypes with different levels of genetic resistance under controlled conditions, Nematropica, 44, 2, 25-30.
 http://journals.fcla.edu/nematropica/article/view/83313
- Das, T.K., Bhattacharyya, R., Sudhishri, S., Sharma, A.R., Saharawat, Y.S., Bandyopadhyay, K.K., Sepat, S.,
 Bana, R.S., Aggarwal, P., Sharma, R.K., Bhatia, A., Singh, G., Datta, S.P., Kar, A., Singh, B., Singh, P.,
 Pathak, H., Vyas, A.K., Jat, M.L. (2014). Conservation agriculture in an irrigated cotton-wheat system

of the western Indo-Gangetic Plains: Crop and water productivity and economic profitability, Field Crops Research, 158, 24-33. <u>http://www.sciencedirect.com/science/article/pii/S0378429013004218</u>

- Dendooven, L., Patiño-Zuñiga, L., Verhulst, N., Boden, K., Garcia-Gaytan, A., Luna-Guido, M., Govaerts, B. (2014.) Greenhouse gas emissions from nontilled, permanent raised, and conventionally tilled beds in the Central Highlands of Mexico, Journal of Crop Improvement, 28, 4, 547-574.
 <u>http://www.tandfonline.com/doi/abs/10.1080/15427528.2014.924328?journalCode=wcim20#.U9k OD_IdUms</u>
- Dias Almeida, G., Nair, S., Borem, A., Cairns, J.E., Trachsel, S., Ribaut, J-M, Banziger, M., Prasanna, B.M., Crossa, J., Babu, R. (2014). Molecular mapping across three populations reveals a QTL hotspot region on chromosome 3 for secondary traits associated with drought tolerance in tropical maize, Molecular Breeding, 34, 2, 701-715. <u>http://link.springer.com/article/10.1007%2Fs11032-014-0068-5</u>
- E. Nhamucho, Mugo, S.N., Kinyua, M., L. Gohole, Tadele Tefera, Mulima, E. (2014). Antibiosis Mechanism of Resistance to Larger Grain Borer, Prostephanus truncatus (Horn)(Coleoptera: Bostrichidae) in Maize, Journal of Entomology, 11, 5, 248-260. <u>http://scialert.net/abstract/?doi=je.2014.248.260</u>
- Edae, E.A., Byrne, P.F., Haley, S.D., Lopes, M.S., Reynolds, M.P. (2014). Genome-wide association mapping of yield and yield components of spring wheat under contrasting moisture regimes, Theoretical and Applied Genetics, 127, 4, 791-807. <u>http://link.springer.com/article/10.1007/s00122-013-2257-8</u>
- Fentaye Kassa Semagn, Beyene, Y., Babu, R., Sudha, K.N., Gowda, M., Das, B., Amsal Tesfaye Tarekegne, Mugo, S.N., Mahuku, G., Regasa, M.W., Warburton, M.L., Olsen, M., Prasanna, B.M. (2014). QTL mapping and molecular breeding for developing stress resilient maize for sub-Saharan Africa, Crop Science, In press.
- Fentaye Kassa Semagn, Magorokosho, C., Ogugo, V., Makumbi, D., Warburton, M.L. (2014). Genetic relationships and structure among open-pollinated maize varieties adapted to eastern and southern Africa using microsatellite markers, Molecular Breeding, 34, 3, 1423-1435. <u>http://link.springer.com/article/10.1007%2Fs11032-014-0126-z</u>
- Fisher, M. (2014). Are Modern Varieties Always Better? An Economic Analysis of Maize Varietal Selection, African Journal of Agricultural and Resource Economics, 9, 4, 270-285. <u>http://www.afjare.org/resources/issues/vol_9_no4/2%20%20Reimer%20%20Fisher.pdf</u>
- Gasura, E., Setimela, P.S., Amsal Tesfaye Tarekegne, Icishahayo, D., Edema, R., Gibson, P.T., Okori, P. (2014). Variability of grain-filling traits in early maturing CIMMYT tropical maize inbred lines, Crop Science, 54, 2, 530-536. <u>https://www.crops.org/publications/cs/abstracts/54/2/530</u>
- Gbegbelegbe, S.D., Chung, U., Shiferaw, B., Msangi, S. (2014). Kindie Tesfaye Fantaye, Quantifying the impact of weather extremes on global food security: A spatial bio-economic approach, Weather and Climate Extremes, 4, 96-108.
 http://www.sciencedirect.com/science/article/pii/S2212094714000474#
- Abu, G., Mekbib, F., Teklewold, A. (2014). Effect of genotype on In vitro propagation of elite sugarcane (Saccharum Officinarum L.) varieties of Ethiopian sugar Estates, International Journal of Technology Enhancements and Emerging Engineering Research, 2, 6, 123-128. <u>http://www.ijteee.org/finalprint/june2014/Effect-Of-Genotype-On-In-Vitro-Propagation-Of-Elite-Sugarcane-saccharum-Officinarum-L-Varieties-Of-Ethiopian-Sugar-Estates.pdf</u>

- Guzman, C., Alvarez, J.B. (2014). Molecular characterization of two novel alleles of Hordoindoline genes in Hordeum chilense Roem. et Schult., Genetic Resources and Crop Evolution, 61, 2, 307-312. http://link.springer.com/article/10.1007%2Fs10722-013-0077-y
- Guzman, C., Medina-Larque, A.S., Velu, G., Gonzalez-Santoyo, H., Singh, R.P., Huerta-Espino, J., Ortiz-Monasterio, I., Peña-Bautista, R.J. (2014). Use of wheat genetic resources to develop biofortified wheat with enhanced grain zinc and iron concentrations and desirable processing quality, Journal of Cereal Science, 60, 3, 617-622.

http://www.sciencedirect.com/science/article/pii/S0733521014001520

- Habash, D. Z., Baudo M., Hindle M., Powers S. J., Defoin-Platel M., Mitchell R., Saqi M., Rawlings C., Latiri K., Araus J. L., Abdulkader A., Tuberosa R., Lawlor D.W. and Nachit M.M. (2014). Systems Responses to Progressive Water Stress in Durum Wheat. PLoS ONE 9(9):e108431.
 DOI:10.1371/journal.pone.0108431.
- Hadgu, G., Kindie Tesfaye Fantaye, Mamo, G.(2014). Analysis of climate change in Northern Ethiopia: implications for agricultural production, Theoretical and Applied Climatology, In press. <u>http://link.springer.com/article/10.1007%2Fs00704-014-1261-5</u>
- Hae Koo Kim, Park, J., Hwang, I.(2014). Investigating water transport through the xylem network in vascular plants, Journal of Experimental Botany, 65, 7, 1895-1904. <u>http://jxb.oxfordjournals.org/content/early/2014/03/07/jxb.eru075</u>
- Hakam, N., Udupa S. M., Gaboun F., Rabha A., Ibriz M. and Iraqi D. (2014). Effect of Genotypes and Culture Media on Embryogenic Callus induction and plantlet Regeneration from Mature Embroyos of Durum Wheat. Romanian Agricultural Research, No. 31, DII 2067-5720 RAR 2014-413.
- He Zhonghu, Xia Xianchun, Peng, S., Lumpkin, T.A.(2014). Meeting demands for increased cereal production in China, Journal of Cereal Science, 59, 3, 235-244. http://www.sciencedirect.com/science/article/pii/S0733521013001501
- Herrera-Foessel, S.A., Huerta-Espino, J., Calvo-Salazar, V., Lan, C.X., Singh, R.P.(2014). Lr72 confers resistance to leaf rust in durum wheat cultivar Atil C2000, Plant Disease, 98, 5, 631-635. <u>http://apsjournals.apsnet.org/doi/abs/10.1094/PDIS-07-13-0741-RE</u>
- Herrera-Foessel, S.A., Singh, R.P., Lillemo, M., Huerta-Espino, J., Bhavani, S., Sukhwinder-Singh, Caixia Lan, Calvo-Salazar, V., Lagudah, E.S. (2014). Lr67/Yr46 confers adult plant resistance to stem rust and powdery mildew in wheat, Theoretical and Applied Genetics, 127, 4, 781-789, <u>http://link.springer.com/article/10.1007/s00122-013-2256-9</u>
- Hickey, J.M., Dreisigacker, S., Crossa, J., Hearne, S., Babu, R., Prasanna, B.M., Grondona, M., Zambelli.
 A., Windhausen, V.S., Mathews, K.L., Gorjanc, G. (2014). Evaluation of genomic selection training population designs and genotyping strategies in plant breeding programs using simulation, Crop Science, 54, 1476-1488. <u>https://www.crops.org/publications/cs/view/first-look/c13-03-0195.pdf</u>
- Karabayev, M., Morgounov, A.I., Braun, H.J., Wall, P.C., Sayre, K.D., Zelenskiy, Y., Zhapayev, R.,
 Akhmetova, A., Dvurechenskii, V., Iskandarova, K., Friedrich, T., Fileccia, T., Guadagnl, M. (2014).
 Effective Approaches to Wheat Improvement in Kazakhstan: breeding and Conservation Agriculture,
 Journal of Agricultural Science and Technology, Accepted for print.
- Karabayev, M., Morgounov, A.I., Wall, P.C., Sayre, K.D., Zelenskiy, Y., Zhapayev, R., Dvurechenskii, V., Akhmetova, A., Friedrich, T., Fileccia, T., GuadagnI, M. (2014). Conservation Agriculture and breeding

for sustainable wheat production in Kazakhstan, Bahri Dagdas Plant Research Journal, Accepted for print.

- Kashiwagi, J., Morito Y., Jitsuyama Y., An P., Inoue T. and Inagaki M. (2014). Effects of root water uptake efficiency on soil water utilization in wheat (Triticum aestivum L.) under severe drought environments. J. Agro. Crop Sci. 200: DOI: 10.1111/jac.12092 (http://onlinelibrary.wiley.com/doi/10.1111/jac.12092/abstract).
- Kassie, G.T., Tsedeke Abate, Langyintuo, A., Maleni, D. (2014). Poverty in maize growing rural communities of, southern Africa, Development Studies Research: An Open Access Journal, 1, 1, 311-323. http://www.tandfonline.com/doi/pdf/10.1080/21665095.2014.969844
- Kher, S.V., Dosdall L. M., Cárcamo H. A.and El-Bouhssini M. (2014). Antixenosis Resistance to Oulema melanopus (Coleoptera: Chrysomelidae) in Central Asian Wheat Germplasm. Journal of Economic Entomology 107: 410-416.
- Khodarahmi, M., Mohammadi, S.A., Bihamta, M.R., Majidi Heravan, E., Jalal Kamali, M.R. (2014). Inheritance and combining ability of yellow rust resistance in some bread wheat commercial cultivars and advanced lines, Seed and Plant Improvement Journal, 30, 3, 531-544. <u>http://spij.spii.ir/browse.php?a_id=750&sid=1&slc_lang=en</u>
- Kindie Tesfaye Fantaye, Jaleta Debello Moti, Jena, P.R., Mutenje, M. (2014). Identifying Potential Recommendation Domains for Conservation, Agriculture in Ethiopia, Kenya, and Mala, Environmental Management, In press. <u>http://link.springer.com/content/pdf/10.1007%2Fs00267-014-0386-8.pdf</u>
- Krishna, V.V., Erenstein, O., Sadashivappa, P., Vivek, B. (2014). Potential Economic Impact of Biofortified Maize in the Indian Poultry Sector, International Food and Agribusiness Management Review, 17, 4, 109-138. <u>http://www.ifama.org/files/IFAMR/Vol%2017/Issue%204/201301025.pdf</u>
- Kukal, S.S., Yadvinder-Singh, Jat, M.L., Sidhu, H.S. (2014). Improving water productivity of wheat-based cropping systems in South Asia for sustained productivity, Advances in Agronomy, 127, 157-258. http://www.sciencedirect.com/science/article/pii/B9780128001318000042
- Lifang Xing, Cuifen Wang, Xianchun Xia, He Zhonghu, Wanquan Chen, Taiguo Liu, Zaifeng Li, Daqun Liu (2014). Molecular mapping of leaf rust resistance gene LrFun in Romanian wheat line Fundulea 900, Molecular Breeding, 33, 4, 931-937. <u>http://link.springer.com/article/10.1007%2Fs11032-013-0007-x</u>
- Linda, M, Nachit M.M. and Augusto P. M. (2014). Allelic variants in durum wheat (Triticum turgidum L. var. durum) DREB genes conferring tolerance to abiotic stresses. Under press in Molecular Genetics and Genomics.
- Loladze, A., Dhouha Kthiri, Pozniak, C., Ammar, K. (2014). Genetic Analysis of Leaf Rust Resistance in Six Durum Wheat Genotypes, Phytopathology, In press. <u>http://apsjournals.apsnet.org/doi/pdfplus/10.1094/PHYTO-03-14-0065-R</u>
- Loladze, A., Druml, T., Wellings, C.R. (2014). Temperature adaptation in Australasian populations of Puccinia striiformis f. sp. tritici, Plant Pathology, 63, 3, 572-580. http://onlinelibrary.wiley.com/doi/10.1111/ppa.12132/abstract
- Long-Xi Yu, Barbier, H., Rouse, M.N., Sukhwinder-Singh, Singh, R.P., Bhavani, S., Huerta-Espino, J., Sorrells, M.E. (2014). A consensus map for Ug99 stem rust resistance loci in wheat , Theoretical and

Applied Genetics, 127, 7, 1561-1581. http://link.springer.com/article/10.1007%2Fs00122-014-2326-7

- Lopez-Vera, E.E., Nelson, S., Singh, R.P., Basnet, B.R., Haley, S.D., Bhavani, S., Huerta-Espino, J., Xoconostle-Cazares, B.G., Ruiz-Medrano, R., Rouse, M.N., Sukhwinder-Singh (2014). Resistance to stem rust Ug99 in six bread wheat cultivars maps to chromosome 6DS, Theoretical and Applied Genetics, 127, 1, 231-239. http://link.springer.com/article/10.1007/s00122-013-2212-8
- Lyimo, S., Mduruma, Z., De Groote, H. (2014). The use of improved maize varieties in Tanzania, African Journal of Agricultural Research, 9, 7, 643-657. <u>http://www.academicjournals.org/article/article1392116190_Lyimo%20et%20al.pdf</u>
- Macharia, G.K., Peña Bautista, R.J., Simsek, S., Anderson, J.A. (2014). Variation at glutenin subunit loci, single kernel characterization and evaluation of grain protein in East African bread wheat varieties, Euphytica, 197, 3, 409-421. <u>http://link.springer.com/article/10.1007%2Fs10681-014-1077-5</u>
- Machida, L., Derera, J., Tongoona, P., Langyintuo, A., MacRobert, J. (2014). Exploration of farmers' preferences and perceptions of maize varieties: implications on development and adoption of quality protein maize (QPM) varieties in Zimbabwe, Journal of Sustainable Development, 7, 2, 194-207. <u>http://www.ccsenet.org/journal/index.php/jsd/article/view/28782</u>
- Maphosa, L., Langridge, P., Taylor, H., Parent, B., Emebiri, L.C., Kuchel, H., Reynolds, M.P., Chalmers, K.J., Okada, A., Edwards, J., Mather, D.E. (2014). Genetic control of grain yield and grain physical characteristics in a bread wheat population grown under a range of environmental conditions, Theoretical and Applied Genetics, 127, 7, 1607-1624.
 http://link.springer.com/article/10.1007%2Fs00122-014-2322-y
- Martinez-Cisneros, B.A., Juarez-Lopez, G., Valencia-Torres, N., Duran-Peralta, E., Mezzalama, M. (2014). First report of bacterial stalk rot of maize caused by Dickeya zeae in Mexico, Plant Disease, 98, 9, 1267. <u>http://apsjournals.apsnet.org/doi/abs/10.1094/PDIS-02-14-0198-PDN</u>
- Mazid, A., Keser, M., Amegbeto, K.N., Morgounov, A.I., Bagci, A., Peker, K., Akin, M., Kucukcongar, M., Kan, M., Semerci, A., Karabak, S., Altikat, A., Yaktubay, S. (2014). Measuring the impact of agricultural research: The case of new wheat varieties in Turkey, Experimental Agriculture, In press. <u>http://journals.cambridge.org/action/displayAbstract?fromPage=online&aid=9317527&fulltextType</u> <u>=RA&fileId=S0014479714000209</u>
- Melloul, M., Iraqi D., Udupa S.M., Elalaoui A., Erba G., Ibriz M. and Elfahime E. (2014). cDNA sequences derived from mRNA induced in leaves in response to water deficit in Triticum turgidum subsp. durum. Submitted NCBI Database. Accession numbers JZ482427-JZ482472.
 http://ebi.edu.au/ftp/databases/dbEST/Daily.FASTA/dbEST.daily.FASTA.082813
- Melloul, M., Iraqi D., Udupa S.M., Erba G., El Alaoui My A., Ibriz M., El Fahime E. (2014). Analysis of mRNA Levels of Ten Genes Under Water Stress in Triticum turgidum subsp. durum. Journal of Plant Studies, 3 (1): DOI: 10.5539/jps.v3n1p65.
- Mergoum, M., Simsek, S., Shaobin Zhong, Acevedo, M., Friesen, T.L., Singh, P.K., Adhikari, T.B., Alamri, M.S., Frohberg, R.C. (2014). 'Velva' spring wheat: an adapted cultivar to North-Central plains of the United States with high agronomic and quality performance, Journal of Plant Registrations, 8, 1, 32-37. <u>https://www.crops.org/publications/jpr/abstracts/8/1/32</u>
- Meti, P., Sreenivas, A.G., Prakash, K., Jat, M.L., Venkateshalu, A., Prabhuraj, N., Manjunath, N., Singh, Y.K. (2014). Population dynamics of shoot fly and stem borers of maize under conservation

agriculture system, Journal of Experimental Zoology India, 17, 2, 563-566. http://www.cabdirect.org/abstracts/20143367858.html

- Mohammadi, R., Haghparast R., Sadeghzadeh B., Ahmadi H., Solimani K.and Amri A. (2014). Adaptation Patterns and Yield Stability of Durum Wheat Landraces to Highland Cold Rainfed Areas of Iran. Crop Sci. 54:944–954.
- Molero, G., Tcherkez, G., Araus, J.L., Nogues, S.(2014). On the relationship between C and N fixation and amino acid synthesis in nodulated alfalfa (Medicago sativa), Functional Plant Biology, 41, 4, 331-341. http://www.publish.csiro.au/nid/102/paper/FP13189.htm
- Morgounov, A., Abugalieva, A., Martynov, A.S.(2014). Effect of climate change and variety on longterm variation of grain yield and quality in winter wheat in Kazakhstan, Cereal Research Communications, 42, 1, 163-172. <u>http://www.akademiai.com/content/y8uu02g84778h231/</u>
- Mulvaney, M.J., Verhulst, N., Herrera, J.M., Mezzalama, M., Govaerts, B. (2014). Improved wheat performance with seed treatments under dry sowing on permanent raised beds, Field Crops Research, 164, 189-198. <u>http://www.sciencedirect.com/science/article/pii/S037842901400118X</u>
- Murungweni, C., Wijk, M.T. van, Giller, K.E., Andersson, J.A., Smaling, E.M.A. (2014). Adaptive livelihood strategies employed by farmers to close the food gap in semi-arid south eastern Zimbabwe, Food Security, 6, 313-326. <u>http://link.springer.com/article/10.1007%2Fs12571-014-0348-2</u>
- Nagel, M., Navakode S., Scheibal V., Baum M., Nachit M., Röder M.S. and Börner A. (2014). The genetic basis of durum wheat germination and seedling growth under osmotic stress. Biologia Plantarum XX (X): XXX-XXX.
- Nazco, R, Peña-Bautista, R.J., Ammar, K., Villegas, D., Crossa, J., Moragues, M., Royo, C. (2014).
 Variability in glutenin subunit composition of Mediterranean durum wheat germplasm and its relationship with gluten strength, Journal of Agricultural Science, 152, 3, 379-393.
 http://cambridgefluids.org/action/displayAbstract?fromPage=online&aid=8851785&fulltextType=R
 http://cambridgefluids.org/action/displayAbstract?fromPage=online&aid=8851785&fulltextType=R
- Nazco, R., Peña Bautista, R.J., Ammar, K., Villegas, D., Crossa, J., Royo, C. (2014). Durum wheat (Triticum durum Desf.) Mediterranean landraces as sources of variability for allelic combinations at Glu-1/Glu-3 loci affecting gluten strength and pasta cooking quality, Genetic Resources and Crop Evolution, 61, 6, 1219-1236. <u>http://link.springer.com/article/10.1007%2Fs10722-014-0104-7</u>
- Ndhlela, T., Herselman, L., Magorokosho, C., Setimela, P.S., Mutimaamba, C., Labuschagne, M. (2014). Genotype X Environment Interaction, of Maize Grain Yield Using AMMI Biplots, Crop Science, 54, 5, 1992-1999. <u>https://www.crops.org/publications/cs/abstracts/54/5/1992</u>
- Nefzaoui, M., Udupa S. M., Gharbi M. S., Bouhadida M. and Iraqi D. (2014). Molecular Diversity in Tunisian Durum Wheat Accessions based on Microsatellite Markers Analysis. Romanian Agricultural Research, No. 31, DII 2067-5720 RAR 2014-359.
- Ngwira, A., Johnsen, F.H., Aune, J.B., Mekuria, M., Thierfelder, C. (2014). Adoption and extent of conservation agriculture practices among smallholder farmers in Malawi, Journal of Soil and Water Conservation, 69, 2, 107-119. <u>http://www.jswconline.org/content/69/2/107.abstract</u>
- Ngwira, A.R., Aune, J.B., Thierfelder, C. (2014). On-farm evaluation of the effects of the principles and components of conservation agriculture on maize yield and weed biomass in Malawi, Experimental

Agriculture, 50, 4, 591-610.

http://journals.cambridge.org/action/displayAbstract?aid=9345087&fileId=S001447971400009X

- Ninou, E., Mylonas I., Tsivelikas A., Ralli P., Dordas Ch., Tokatlidis I. (2014). Wheat landraces are better qualified as potential gene pools at ultraspaced rather than densely grown conditions. The Scientific World Journal. http://dx.doi.org/10.1155/2014/957472.
- Ogugo, V., Semagn, K., Beyene, Y., Runo, S., Olsen, M., Warburton, M.L. (2014). Parental genome contribution in maize DH lines derived from six backcross populations using genotyping by sequencing, Euphytica, In press. <u>http://link.springer.com/article/10.1007/s10681-014-1238-6</u>
- Ortega, R., Alvarez, J.B., Guzman, C. (2014). Characterization of the Wx gene in diploid Aegilops species, and its potential use in wheat breeding, Genetic Resources and Crop Evolution, 61, 2, 369-382. <u>http://link.springer.com/article/10.1007%2Fs10722-013-0040-y</u>
- Ortega, R., Guzman, C., Alvarez, J.B. (2014). Wx gene in diploid wheat: molecular characterization of five novel alleles from einkorn (Triticum monococcum L. ssp. monococcum) and T. urartu, Molecular Breeding, 34, 3, 1137-1146. <u>http://link.springer.com/article/10.1007%2Fs11032-014-0105-4</u>
- Pariyar, S.R., Dababat, A.A., Nicol, J.M., Orakci, G.E., Goll, M.B., Watrin, C., Duveiller, E., Braun, H.J., Sikora, R. (2014). Fungicide seed treatment and host resistance for the management of wheat crown rot caused by Fusarium, culmorum, Basic Research Journal of Agricultural Science and Review, 3, 9, 116-121. http://www.basicresearchjournals.org/agric/abstract/Pariyar%20et%20al.html
- Patpour, M., Nazari K., Ogbonnaya F., Alavi S.M. and Mousavi A. (2014). Phenotypic and Molecular Characterization of Resistance to Stem Rust in Wheat Cultivars and Breeding Advanced Lines from Iran and Syria. Crop Breeding Journal (accepted for Volume 3, Number 2, 2013 (in press).
- Rehman, A., Sajjad, M., Khan, S.H., Peña Bautista, R.J., Khan, N.I. (2014). Lower tendency of allelic variation of Glu genes and absence of 1BL-1RS translocation in modern Pakistani wheats, Cereal Research Communications, 42, 1, 139-150.
 http://www.akademiai.com/content/22550n1n111w6n48/?p=ad4421acadcc4e62a87ef7a756a4a98 9&pi=5
- Roostaei, M., Mohammadi R. and Amri A. (2014). Rank correlation among different statistical models in ranking of winter wheat genotypes. The Crop Journal 2:154-163.
- Sanchez-Garcia, M., Álvaro F., Peremarti A., Trevaskis B., Martín-Sánchez J. A. and Royo C. (2014). Breeding effects on dry matter accumulation and partitioning in Spanish bread wheat during the 20th century. Euphytica DOI:10.1007/s10681-014-1268-0.
- Sharma, P.C., Singh, D., Sehgal, D., Singh, G., Hash, C.T., Yadav, R.S. (2014). Further evidence that a terminal drought tolerance QTL of pearl millet is associated with reduced salt uptake, Environmental and Experimental Botany, 102, 48-57. http://www.sciencedirect.com/science/article/pii/S0098847214000392
- Sharma, R.C., Morgounov A., Akin B., Bespalova L., Lang L., Litvinenko M., Mustatea P., Ozturk I.,
 Postolatiy A., Rajaram S. and Braun H.J. (2014). Winter wheat East European Regional Yield Trial:
 Identification of superior genotypes and characterization of environments. Crop Science 54:2469-2480.

- Shujun Wang, Heyang Luo, Jian Zhang, Yan Zhang, He Zhonghu, Shuo Wang, (2014). Alkali-induced changes in functional properties and in vitro digestibility of wheat starch: the role of surface proteins and lipids, Journal of Agricultural and Food Chemistry, 62, 16, 3636-3643. http://pubs.acs.org/doi/abs/10.1021/jf500249w
- Snapp, S.S., Fisher, M. (2014). "Filling the maize basket" supports crop diversity and quality of household diet in Malawi, Food Security, In press. <u>http://link.springer.com/article/10.1007/s12571-014-0410-0</u>
- Sohail, Q., Shehzad T., Kilian A., Eltayeb A.E., Tanaka H. and Tsujimoto H. (2012). Development of diversity array technology (DArT) markers for assessment of population structure and diversity in Aegilops tauschii. Breed Sci. 62: 38-45.
- Soleiman, N.H., Solis, I., Ammar, K., Dreisigacker, S., Soleiman, M.H., Martinez, F., Loladze, A. (2014). Resistance to leaf rust in a set of durum wheat cultivars and landraces in Spain, Journal of Plant Pathology, 96, 2, 353-362.
- Soleiman, N.H., Solis, I., Sillero, J.C., Herrera-Foessel, S.A., Ammar, K., Martinez, F. (2014). Evaluation of macroscopic and microscopic components of partial resistance to leaf rust in durum wheat, Journal of Phytopathology, 162, 6, 359-366. <u>http://onlinelibrary.wiley.com/doi/10.1111/jph.12197/abstract</u>
- Srinivasa, J., Arun, B., Mishra, V.K., Chand, R., Sharma, D., Bhardwaj, S.C., Joshi, A.K. (2014). Accessing the spelt gene pool to develop well-adapted bread wheat lines with increased grain zinc and iron, Crop Science, 54, 5, 1-11. <u>https://www.crops.org/publications/cs/view/first-look/c13-12-0801.pdf</u>
- Sukumaran, S., Jianming Yu, Association mapping of genetic resources: Achievements and future perspectives, 2014, Genomics of Plant Genetic Resources, p. 207-235.
- Tadesse, W., Nachit M., Abdalla O. and Rajaram S. (2014). Wheat Breeding at ICARDA: Achievements and Prospects in the CWANA Region. In Alain Bonjean, Bill Angus and Maarten van Ginkel (eds). World Wheat BooK Volume 3.
- Tadesse, W., Ogbonnaya F. C., Jighly A., Sanchez-Garcia M., Sohail Q., Rajaram S. and Baum M. (2014).
 Genome-wide Association Mapping of Yield and Grain Quality Traits in Elite Winter Wheat
 Genotypes Targeted to the CWANA region. Mol. Breeding (under review).
- Tadesse, W., Ogbonnaya F.C., Jighly A., Nazari K., Rajaram S. and Baum M. (2014). Association Mapping of Resistance to Yellow Rust in Winter Wheat Cultivars and Elite Genotypes. Crop science, vol. 54. Joukhadar, R., El-Bouhssini M., Jighly A. and Ogbonnaya F. C. (2014). Genome-wide association mapping for five major pest resistances in wheat.Springer, Mol Breeding, DOI 10.1007/s11032-013-9924-y. Park, R. F., Mohler V., Nazari K. and Singh D. (2014). Characterisation and mapping of gene Lr73 conferring seedling resistance to Puccinia triticina in common wheat. Theor Appl Genet (2014) 127:2041–2049. DOI 10.1007/s00122-014-2359-y.
- Tadesse, W., Rajaram S., Ogbonnaya F. C., Sanchez-Garcia M., Sohail Q. and Baum M. (2014). Wheat Breeding: Challenges and Prospects in the 21st Century. Springer.
- Tinak Ekom, D.C., Udupa S. M., Benchekroun M. N., Ennaji M. M., Abdelwahd R.and Iraqi D. (2014). Immature Embryo-Derived of Two Bread Wheat (Triticum aestivum L.) Varieties Transformation Using Particle Bombardment Method. Annual Research & Review in Biology, 4(24): 3904-3914.

- Turner, M.D., Butt, B., Singh, A., Brottem, L., Ayantunde, A., Gerard, B. (2014). Variation in vegetation cover and livestock mobility needs in Sahelian, West Africa, Journal of Land Use Science, In press. http://www.tandfonline.com/doi/full/10.1080/1747423X.2014.965280#.VG4vk4vF-X0
- Vargas, M., Glaz, B., Alvarado, G., Pietragalla, J., Morgounov, A.I., Zelenskiy, Y., Crossa, J. (2014). Analysis and interpretation of interactions in agricultural research, Agronomy Journal, In press. https://www.agronomy.org/publications/aj/view/first-look/aj13-0405.pdf
- Velu, G., Ortiz-Monasterio, I., Cakmak, I., Yuanfeng Hao, Singh, R.P. (2014). Biofortification strategies to increase grain zinc and iron concentrations in wheat, Journal of Cereal Science, 59, 3, 365-372. <u>http://www.sciencedirect.com/science/article/pii/S0733521013001495#</u>
- Yildirim, M., Gezginc, H., Paksoy, A.H., Aktas, B., Cakmak, M., Belen, S., Bolat, N., Yorgancilar, O., Keser, M., Braun, H.J., Pfeiffer, W.H. (2014). A study on some characteristics of new registered triticale (x Triticosecale Wittm.) varieties (Aysehanim and Mehmetbey) in Turkey, Journal of Food, Agriculture and Environment, 12, 2, 370-374. <u>http://world-food.net/a-study-on-some-characteristics-of-newregistered-triticale-x-triticosecale-wittm-varieties-aysehanim-and-mehmetbey-in-turkey/</u>
- Yonglu Tang, Rosewarne, G.M., Chaosu Li, Xiaoli Wu, Wuyun Yang, Chun Wu (2014). Physiological factors underpinning grain yield improvements of synthetic derived wheat in South Western China, Crop Science, In press. <u>https://www.crops.org/publications/cs/view/first-look/crop-2014-02-0124-ora.pdf</u>
- Zaifeng Li, Caixia Lan, He Zhonghu, Singh, R.P., Rosewarne, G.M., Xinmin Chen, Xianchun Xia (2014). Overview and Application of QTL for Adult Plant Resistance to Leaf Rust and Powdery Mildew in Wheat, Crop Science, 54, 5, 1907-1925. https://www.crops.org/publications/cs/abstracts/54/5/1907
- Zarei, B., Naderi, A., Jalal Kamali, M.R., Lack, S., Modhej, A. (2014).Phonological traits, yield components and their relationship with grain yield stability in spring wheat genotypes under terminal heat stress conditions, Advances in Environmental Biology, 8, 10, 1177-1184.
 http://www.academia.edu/8227780/Advances in Environmental Biology Phonological Traits Yiel d Components and their Relationship with Grain Yield Stability in Spring Wheat Genotypes Under Terminal Heat Stress Conditions INTRODUCTION
- Zhang, X., Perez-Rodriguez, P., Fentaye Kassa Semagn, Beyene, Y., Babu, R., Lopez-Cruz, M., San Vicente, F.M., Olsen, M., Buckler, E., Jannink, J-L., Prasanna, B.M., Crossa, J. (2014). Genomic prediction in biparental tropical maize populations in water-stressed and well-watered environments using low-density and GBS SNPs, Heredity, In press. http://www.nature.com/hdy/journal/vaop/ncurrent/full/hdy201499a.html
- Zhapayev, R., Iskandarova, K., Toederich, K., Paramonova, I., Andybayev, D., Al-Dakheel, A., Ismail, S., Pinnamanent, S., Nekrasova, N., Balpanov, D., Ten, O., Ramanculov, E., Zelenskiy, Y., Karabayev, M., Akhmetova, A. (2014). Sweet sorghum genotypes testing in the high latitude rainfed steppes of the North Kazakhstan, Journal of Arid Lands Studies, Accepted for print.

Conferences, Workshops and Symposia

Afshari, F., Nazari K., Aghaei M., Najafian G., Esmaeilzadeh M., Yazdansepas A., Khodarahmei M., Amini A., Roohparvar R., Hashamei M., Malihipour A. and Patpour M. (2014). The wheat stripe rust pathogen in Iran and achievement of resistant wheat cultivars over the last 10 years. 2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.

- Amri, A., El Bouhssini M., Nsarellah N., Nachit M., Abdallah O., Lhaloui S., Jlibene M., Mergoum M., El Haddoury J., ElHaila M., Gill B., Cox S.and Hatchett J. (2014). Results of collaborative germplasm enhancement for resistance to Hessian fly of wheat in Morocco. International Plant Resistance to Insects Workshop, Marrakech, Morocco, 14-16 April, 2014.
- Bari, A., Nazari K., Nachit M., Amri A., Street K., Yahyaoui A. and Endresen D. (2014). Predicting and locating sources of resistance to stripe (yellow) rust in durum wheat genetic resources. 2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.
- Cherif, A., El Bouhssini M., Glida H., Gharbi M.S. and Mediouni Ben Jemâa J. (2014). Potential of genes H22, H18, H13 and H11 for durum wheat resistance to Mayetiola destructor field populations from North Tunisia. International Plant Resistance to Insects Workshop, Marrakech, Morocco, 14-16 April, 2014.
- Dusunceli, F., ElKhoury W., Mancini F., Hodson D., Nazari K., Ramdani A., Muminjanov H. and Nersisyan A. (2014). Guidelines for development of comprehensive national policies for integrated management of the wheat rusts. Proceedings of BGRI 2014 Technical Workshop 22-25 March Obregon, Mexico.
- Dusunceli, F., Khoury W., Muminjanov H., Nersisyan A., Aldobai S., Birisik N., Sahin M., Selcuk S., Asikoglu T., Mert Z., Akan K., Nazari K. and Hodson D. (2014). Experiences and vision of FAO on surveillance and management of wheat rusts gained through its Wheat Rust Diseases Global Program. 2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.
- El Amil, R., Leconte M., Hovmøller M., Shykoff J., Nazari K. and Vallavieille-Pope C.de. (2014). Effect of the temperature on aggressiveness components, infection efficiency and latency period, on Puccinia striiformis f. sp. tritici in the Middle East..2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.
- El-Hanafi, S., Henkrar F., El-Haddoury J., Bendaou N.and Sripada S. (2014). Improving a Moroccan wheat cultivar 'Aguilal' for resistance to leaf and yellow rust and quality. 2nd International Conference of Plant Biodiversity, PP 151.
- Hansen J. G., Nazari K., Hodson D., Lassen P. and Hovmøller M. S. (2014). Early warning for yellow rust tools and services provided by the wheat rust toolbox. 2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.
- Henkrar, F., El-Haddoury J., Bendaou N. and Udupa S. (2014). Determination of allelic variation at HMW-GS and LMW-GS loci in Moroccan bread and durum wheat cultivars using gene-specific PCR markers. 2nd International Conference of Plant Biodiversity, PP 43.
- Henkrar, F., El-Hanafi S., El-Haddoury J., Bendaou N. and Udupa S. (2014). Anther culture and markerassisted fast track improvement of bread wheat for rust resistance and bread making quality. 2nd International Conference of Plant Biodiversity, PP 129.
- Keser, M., Morgounov A., Akın B., Mert Z., Ozdemir F. and Nazari K. (2014). Developing yellow rust resistant winter wheat germplasm for CWANA. 2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.

- Lhaloui, S., El Bouhssini M., Ouabbou H., Ogbonnaya F., Parry M., Udupa S., Nsarellah N., El Haddouri J. and El Fakhouri K. (2014). New sources of resistance to the Moroccan populations of Hessian fly identified in primary synthetic and reverse genetic strategy-induced point mutation bread wheat lines. International Plant Resistance to Insects Workshop, Marrakech, Morocco, 14-16 April, 2014.
- Morgounov, A., Akan K., Akın B., Belen S., Çetin L., Keser M., Kan M., Küçükçongar M., Mert N.Z., Muminjanov H., Nazari K., Özdemir F., Özer E., Özseven İ. and Qualset C. (2014). Resistance to yellow rust in bread wheat landraces currently grown in Turkey. 2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.
- Moukahel, A.R., Nazari K., S. Asaad, Debbes B., Morris C.E. and Sands D.C. (2014). The impact of the icenucleation bacteria on wheat stripe rust responses under greenhouse conditions. 2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.
- Nazari, K. (2014). Wheat stripe rust surveys in CWANA and a Regional Cereal Rust Research Center at Izmir, Turkey. Proceedings of BGRI 2014 Technical Workshop 22-25 March Obregon, Mexico.
- Nazari, K., Mohler V. and Wellings C. (2014). QTL mapping of resistance to wheat stripe rust in a Cappelle Desprez × Avocet 'S' doubled haploid wheat population. 2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.
- Nazari, K., Sari A.O. and Keser M. (2014). Turkey-ICARDA Rust Collaborative Project: Roadmap of the Regional Cereal Rust Research Center. 2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.
- Nsarellah, N., Nachit M., Lhaloui S., Amri A. and El Bouhssini M. (2014). Recent durum wheat variety release with Hessian fly resistance from inter specific crosses in Morocco. International Plant Resistance to Insects Workshop, Marrakech, Morocco, 14-16 April, 2014.
- Ramdani, A. and Nazari K. (2014). Effective Yr genes under Moroccan conditions during the 2011-2012 and 2012- 2013 cropping seasons. 2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.
- Ramdani, A., Nazari K. and Lhaloui S. (2014). Status of wheat rust diseases in Morocco during the 2012-2013 cropping season. 2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.
- Sharma, R.C., Amanov A., Ziyadullaev Z., Saidov S., Ahmedov M., Bedoshvili D., Kokhmetova A., Keser M., Morgounov A., Nazari K., Rajaram S. and Baum M. (2014). Status of stripe rust resistant winter wheat varieties in Central Asia and the Caucasus. 2nd International Wheat Stripe Rust Symposium, April 28 to May1st, 2014, Izmir, Turkey.
- Tadesse, W. (2014). Wheat Improvement for Dry land Agriculture.In: 53rd All India Wheat and Barley Meet, 22-25 August, Jabalpur, India.
- Tadesse, W., Abdalla O., Rajaram S., Ogbonnaya F., Hagras A., Keser M., Sharma R., Nazari K., Sanchez Garcia M., Sohail Q., Al-Abdallat A. and Baum M. (2014). Bread wheat breeding for yellow rust

resistance at ICARDA: Current status and prospects. Second International Yellow rust Conference, April 28-May1, Izmir, Turkey.

Tadesse, W., Ogbonnaya F. C., Abdella O., El Bouhssini M., Lhaloui S., Ammar A. and Baum M. (2014).
 Identification and development of elite bread wheat genotypes with high yield potential and
 Hessian fly resistance. International Plant Resistance to Insects work shop, April 14-17, Marrakech,
 Morocco.